

ORB5: A global electromagnetic gyrokinetic code using the PIC approach in toroidal geometry[☆]

E. Lanti^{a,*}, N. Ohana^a, N. Tronko^b, T. Hayward-Schneider^b, A. Bottino^b, B.F. McMillan^d, A. Mishchenko^c, A. Scheinberg^e, A. Biancalani^b, P. Angelino^f, S. Brunner^a, J. Dominski^e, P. Donnel^a, C. Gheller^a, R. Hatzky^b, A. Jocksch^g, S. Joliet^a, Z.X. Lu^b, J.P. Martin Collar^b, I. Novikau^b, E. Sonnendrücker^b, T. Vernay^a, L. Villard^a

^a École Polytechnique Fédérale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015 Lausanne, Switzerland

^b Max-Planck-Institut für Plasmaphysik, D-85748 Garching, Germany

^c Max-Planck-Institut für Plasmaphysik, D-17491 Greifswald, Germany

^d CFSA, Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

^e Princeton Plasma Physics Laboratory, Princeton, NJ 08540, USA

^f Swiss Institute of Bioinformatics (SIB), Lausanne, Switzerland

^g CSCS, Swiss National Supercomputing Centre, Via Trevano 131, 6900 Lugano, Switzerland

ARTICLE INFO

Article history:

Received 6 May 2019

Received in revised form 17 September 2019

Accepted 20 November 2019

Available online 30 November 2019

Keywords:

Tokamak
Gyrokinetic
PIC
Turbulence

ABSTRACT

This paper presents the current state of the global gyrokinetic code ORB5 as an update of the previous reference (Joliet et al., 2007). The ORB5 code solves the electromagnetic Vlasov–Maxwell system of equations using a PIC scheme and also includes collisions and strong flows. The code assumes multiple gyrokinetic ion species at all wavelengths for the polarization density and drift-kinetic electrons. Variants of the physical model can be selected for electrons such as assuming an adiabatic response or a “hybrid” model in which passing electrons are assumed adiabatic and trapped electrons are drift-kinetic. A Fourier filter as well as various control variates and noise reduction techniques enable simulations with good signal-to-noise ratios at a limited numerical cost. They are completed with different momentum and zonal flow-conserving heat sources allowing for temperature-gradient and flux-driven simulations. The code, which runs on both CPUs and GPUs, is well benchmarked against other similar codes and analytical predictions, and shows good scalability up to thousands of nodes.

© 2019 EUROfusion. Published by Elsevier B.V. All rights reserved.

1. Introduction

Understanding the critical phenomena limiting the performance of magnetic confinement devices is crucial to achieve a commercially viable fusion energy production. Among them, microinstabilities play a key role as they are closely linked to the tokamak confinement properties. For example, turbulent transport induced by microinstabilities mainly governs the heat and particle losses in toroidally confined plasmas. Another important issue is the interaction between waves and energetic particles produced by the fusion process or resulting from the application of heating by neutral beam injection (NBI) or ion cyclotron range of frequencies (ICRF). In this case, the energetic particles interact with the bulk plasma and destabilize various eigenmodes of the shear Alfvén wave such as toroidal Alfvén eigenmodes (TAE)

or the energetic particle modes (EPM), which deteriorate the confinement properties.

It is shown both experimentally [1,2] and theoretically [3] that these drift-wave-type microinstabilities as well as Alfvén eigenmodes [4,5] have a low frequency compared to the ion gyro frequency of strongly magnetized plasmas and are of small amplitude in the core region compared to the background quantities. This motivates the use of gyrokinetic theory [6,7] which retains a kinetic description of the problem while reducing the numerical cost for solving the equations by removing the fast gyro angle dependence of the system in a consistent way and thus reducing the phase space dimensionality from 6D to 5D.

Among the three main numerical approaches used to solve the gyrokinetic equations [8]: Lagrangian [9–15], Eulerian [16–21], and Semi-Lagrangian [22], the Lagrangian particle-in-cell (PIC) scheme [23] was the first introduced in the context of gyrokinetic simulations [24]. It consists of initially sampling the phase space using numerical particles, also called markers, that represent a portion of the phase space and following their orbit in the 5D space.

[☆] The review of this paper was arranged by Prof. David W. Walker.

* Corresponding author.

E-mail address: emmanuel.lanti@epfl.ch (E. Lanti).

The ORB5 code is a nonlinear global PIC code used for solving the gyrokinetic Vlasov–Maxwell system accounting for the presence of collisions and sources. It is a “full- F ” code, in spite of the δf splitting, which is used as a control variate. No scale separation and no ordering assumption are made on the full distribution function, with the exception that the quasineutrality and Ampère’s equations are linearized about an equilibrium distribution function. The ORB5 code is based on a 3D finite element representation of the fields using B-spline basis functions up to third order. It uses toroidal magnetic coordinates and a field-aligned Fourier filter which drastically reduces particle noise. Originally presented in [25] and further developed in [9] for the electrostatic (ES) and collisionless limit, the ORB5 code has since undergone a substantial amount of additions. Those improvements are targeting the physical models, with e.g. drift-kinetic electron dynamics, electromagnetic (EM) perturbations [26], multiple gyrokinetic ion species, inter and intraspecies collisions [27], hybrid electron model [28,29], removal of the long wavelength approximation [30], various heating sources [31,32] and strong flows [33], and the numerical side with e.g. the enhanced control variate [34–36], and, more recently, the mixed-representation “pullback” scheme [37] resolving the so-called cancellation problem for EM simulations, various noise control operators (generalized moment-conserving Krook operator [31], coarse graining [38], and quadtree [39]), and a thorough refactoring with multithreading using OpenMP and OpenACC which will be detailed in a separate publication. The ORB5 code has been carefully and extensively benchmarked against various Lagrangian, Eulerian, and semi-Lagrangian gyrokinetic codes and always showed a good agreement in the results, see for example [29,30,40–49]. The aim of this paper is to review these improvements, and present the current status of the code and illustrate its performance and capabilities with a few significant results.

The present paper is organized as follows. Section 2 presents the gyrokinetic model implemented in ORB5. Section 3 describes the numerical implementation of the gyrokinetic equations as well as the numerical methods used in the code. The parallel efficiency and a few illustrative, physically relevant simulation results are presented in Section 4. Finally, Section 5 presents the conclusions and future work.

2. Gyrokinetic model

2.1. Magnetic geometry, coordinate system, and normalization

The background fields of a tokamak are usually approximated as axisymmetric. A general axisymmetric magnetic field in the nested-flux-surface region may be expressed as

$$\mathbf{B} = F(\psi)\nabla\varphi + \nabla\psi \times \nabla\varphi, \quad (1)$$

where $F(\psi)$ is the poloidal current flux function, ψ is the poloidal magnetic flux and φ is the toroidal angle. The ORB5 code uses ideal-MHD equilibria, solutions of the Grad–Shafranov equation, that are provided by the CHEASE code [50]. It can also use an analytical *ad-hoc* magnetic equilibrium comprising circular concentric magnetic surfaces.

A straight-field-line coordinate system is used in ORB5. The magnetic surfaces are labeled by $s = \sqrt{\psi/\psi_{\text{edge}}}$ where ψ_{edge} is the value of ψ at the radial edge, the toroidal angle is φ , and the straight-field-line poloidal angle is defined by

$$\theta^* = \frac{1}{q(s)} \int_0^\theta \frac{\mathbf{B} \cdot \nabla\varphi}{\mathbf{B} \cdot \nabla\theta'} d\theta', \quad (2)$$

where $q(s)$ is the safety factor profile and θ is the geometric poloidal angle.

All the physical quantities in ORB5 are normalized according to four reference parameters; these normalizations are used internally and in the code output. The reference quantities are the ion mass m_i , the ion charge $q_i = eZ_i$ with e being the elementary charge and Z_i the ion atomic number, the magnetic field amplitude at the magnetic axis B_0 , and the electron temperature $T_e(s_0)$ at a reference magnetic surface s_0 . Note that for simulations with multiple ion species, the user must define a reference ion species for the normalization. Derived units are then defined with respect to these four parameters: time is in units of the inverse of the ion cyclotron frequency $\Omega_{ci} = q_i B_0 / m_i c$ with c the speed of light in vacuum (CGS units are used in this paper), velocities are normalized to the ion sound velocity $c_s = \sqrt{eT_e(s_0)/m_i}$, lengths are given in units of the ion sound Larmor radius $\rho_s = c_s / \Omega_{ci}$, and the densities are normalized to the volume averaged density $\bar{n}$. These reference quantities are then used to construct normalizations for other quantities in the code such as the electrostatic potential, various fluxes, etc.

2.2. GK equations for fields and particles

The gyrokinetic Vlasov–Maxwell model implemented in ORB5 is derived from a variational principle [51,52] which has some advantages compared to models implemented in other gyrokinetic codes which are derived outside of such a framework. The first advantage is the possibility to include all necessary approximations into the expression of the action before deriving the equations of motion. The second advantage consists of the possibility to consistently derive exactly conserved quantities, corresponding to the model, such as the energy. In the ORB5 code these quantities are then used for diagnostics and the verification of the quality of the simulations. Finally, the variational formulation directly provides the weak form of gyrokinetic Poisson and Ampère equations suitable for a finite element discretization.

The choice of the ordering plays a crucial role in defining the complexity of the gyrokinetic model, and in particular the nonlinear terms which are taken into account. The gyrokinetic variational principle corresponding to the ORB5 model is established according to the specific gyrokinetic ordering achievable for numerical implementation. In particular, it means that all the geometrical effects due to the non-uniformity of the background magnetic field are considered one order smaller than the relative fluctuations of the electromagnetic fields. To quantify that statement, we define the magnetic field geometry-related small parameter $\epsilon_B = \rho_{th}/L_B$, where ρ_{th} is the thermal Larmor radius of the particle and $L_B = |\nabla B/B|^{-1}$ sets up the length scale of the background magnetic field variation. The electromagnetic-fluctuations-related small parameter is defined by $\epsilon_\delta \sim |\mathbf{B}_1|/B \sim c|\mathbf{E}_{1\perp}|/(Bv_{th}) \sim (k_\perp \rho_{th}) e\phi_1/T_i \equiv \epsilon_\perp e\phi_1/T_i$, where $\mathbf{E}$ is the electric field, v_{th} is the thermal velocity, $k_\perp$ is the wave number perpendicular to the magnetic field, B the amplitude of the background magnetic field, ϕ_1 is the perturbed electrostatic potential, T_i is the ion temperature, the subscript 1 refers to the fluctuating part of the corresponding fields and the subscript $\perp$ represents the component perpendicular to the magnetic field line. The parameter $\epsilon_\perp$ allows the distinction between the gyrokinetic theory with $\epsilon_\perp \sim \mathcal{O}(1)$ and the drift-kinetic theory with $\epsilon_\perp \ll 1$. Both models are implemented in the code ORB5.

As shown in [52,53] the ordering $\epsilon_B = \epsilon_\delta^2$ corresponds to gyrokinetic models implemented in most global codes. In particular, it has been demonstrated that the ORB5 equations can be derived via variational calculation from the field-particle Lagrangian accurate to second order with respect to the parameter ϵ_δ . Below we present the variational framework and summarize the main gyrokinetic equations resulting from the variational derivation.

The expression of the action functional leading to the ORB5 code gyrokinetic Maxwell–Vlasov equations containing first order

geometric corrections, i.e. $\mathcal{O}(\epsilon_B)$ terms, and the electromagnetic corrections up to the second order, i.e. $\mathcal{O}(\epsilon_\delta^2)$ terms, is given by:

$$\begin{aligned} \mathcal{A} = \int_{t_0}^{t_1} dt \mathcal{L} = \sum_{\sigma} \int dt d\Omega \left(\frac{q_{\sigma}}{c} \mathbf{A}^* \cdot \dot{\mathbf{X}} \right. \\ \left. + \frac{m_{\sigma} c}{q_{\sigma}} \mu \dot{\Theta} - H_0 \right) f_{\sigma} \\ - \epsilon_{\delta} \sum_{\sigma \neq e} \int dt d\Omega H_1 f_{\sigma} - \epsilon_{\delta} \int dt d\Omega H_1^{\text{dk}} f_e \\ - \epsilon_{\delta}^2 \sum_{\sigma \neq e} \int dt d\Omega H_2 f_{\text{eq},\sigma} \\ - \alpha \epsilon_{\delta}^2 \int dt d\Omega H_2^{\text{dk}} f_{\text{eq},e} \\ - \alpha \epsilon_{\delta}^2 \int dt dV \frac{|\nabla_{\perp} A_{1\parallel}|^2}{8\pi}, \end{aligned} \quad (3)$$

where $\alpha = 0$ corresponds to the electrostatic model and $\alpha = 1$ to the electromagnetic model, $d\Omega = dV dW$ with $dV = d^3\mathbf{X}$ and $dW = B_{\parallel}^* d\mu dp_z$ represents the infinitesimal volume of the reduced (gyrocenter) phase space, $B_{\parallel}^*$ is defined as the parallel component of the symplectic magnetic field $\mathbf{B}^* = \nabla \times \mathbf{A}^*$ with $\mathbf{A}^* = \mathbf{A} + (c/q_{\sigma}) p_z \hat{\mathbf{b}}$ being the symplectic magnetic potential and $\hat{\mathbf{b}}$ being the unit vector parallel to the unperturbed magnetic field line. The action is derived using the p_z formulation in which we define the reduced gyrocenter position $\mathbf{X}$, the canonical gyrocenter momentum $p_z = m_{\sigma} v_{\parallel} + \alpha \epsilon_{\delta} (q_{\sigma}/c) A_{1\parallel}$ with the parallel velocity $v_{\parallel}$, the magnetic moment μ and the fast gyroangle Θ . The sums are made over all the species σ except for the second and third sums where the electrons are excluded because they are treated as drift-kinetic. The first and the second terms of the gyrokinetic action are gyrocenter contributions and the last term is a contribution from the perturbed magnetic field.

Before presenting the equations of motion implemented in ORB5, we discuss all necessary approximations included in the gyrokinetic action given by Eq. (3). The first three terms of the action involve the full distribution functions f_{σ} , while the fourth and fifth terms, involving the nonlinear Hamiltonian H_2 , involve equilibrium distribution functions $f_{\text{eq},\sigma}$, which are by definition invariant under the unperturbed Hamiltonian dynamics, i.e. they satisfy the condition $\{f_{\text{eq},\sigma}, H_0\} = 0$. This approximation brings several simplifications in the model. First, it results in the linearization of the gyrokinetic Poisson and Ampère equations. Second, it simplifies the gyrokinetic Vlasov equation by excluding some nonlinear terms from the gyrocenter characteristics associated with the Hamiltonian H_2 .

The gyrocenter model is fixed via the Hamiltonian terms H_0 (non-perturbed dynamics), H_1 (linear gyrocenter dynamics), H_1^{dk} (linear drift-kinetic dynamics for electrons), and H_2 (nonlinear second order gyrocenter dynamics). The choice of the linear H_1 , H_1^{dk} and nonlinear Hamiltonians H_2 determines the expressions for the gyrokinetic charge and current in the reduced Poisson and Ampère equations. In this section we present the general electromagnetic model of the ORB5 code. For further options and approximations implemented on the level of the reduced particle dynamics, see the sections below.

Concerning the field part of the gyrokinetic action, three approximations have been made. First of all, the quasi-neutrality approximation, which allows one to neglect the perturbed electric field energy $-\epsilon_{\delta}^2 \int dt dV |\mathbf{E}_1|^2 / 8\pi$. The second approximation consists of neglecting the magnetic compressibility of perturbations, i.e. the parallel component of the perturbed magnetic field $B_{1\parallel} = \epsilon_{\delta} |\mathbf{B}_{1\perp}|$ is neglected and only the perpendicular part of the perturbed magnetic field $\mathbf{B}_{1\perp} = \hat{\mathbf{b}} \times \nabla A_{1\parallel}$, associated with

$A_{1\parallel}$, is implemented. Finally, due to the chosen ordering, the background component of the magnetic field can be excluded from the Maxwell part of the gyrokinetic action.

The background Hamiltonian contains information about the kinetic energy of a charged particle moving in the unperturbed magnetic field with amplitude B :

$$H_0 = \frac{p_z^2}{2m_{\sigma}} + \mu B. \quad (4)$$

The linearized Hamiltonian model for ions is given by the gyroaveraged linear electromagnetic potential:

$$H_1 = q_{\sigma} \left\langle \phi_1 - \alpha A_{1\parallel} \frac{p_z}{m_{\sigma} c} \right\rangle, \quad (5)$$

where $\langle \dots \rangle$ is the gyroaveraging operator. The gyroaveraging is absent in the linear Hamiltonian model for the electrons because they are considered as drift-kinetic:

$$H_1^{\text{dk}} = -e \left(\phi_1(\mathbf{X}) - \alpha A_{1\parallel}(\mathbf{X}) \frac{p_z}{m_e c} \right). \quad (6)$$

The nonlinear Hamiltonian model which contains all orders in finite Larmor radius (FLR) in its electrostatic part and up to second order FLR terms in its electromagnetic part is considered for ions only:

$$\begin{aligned} H_2 = -\frac{q_{\sigma}^2}{2B} \frac{\partial}{\partial \mu} \left\langle \tilde{\phi}_1 (\mathbf{X} + \boldsymbol{\rho}_0)^2 \right\rangle \\ + \alpha \frac{q_{\sigma}^2}{2m_{\sigma} c^2} \left[A_{1\parallel}(\mathbf{X})^2 + m_{\sigma} \left(\frac{c}{q_{\sigma}} \right)^2 \frac{\mu}{B} A_{1\parallel}(\mathbf{X}) \nabla_{\perp}^2 A_{1\parallel}(\mathbf{X}) \right], \end{aligned} \quad (7)$$

where $\tilde{\phi}_1$ represents the fluctuating part of a perturbed electrostatic potential and $\boldsymbol{\rho}_0$ is the lowest order guiding-center displacement. Finally the second order Hamiltonian for the electrons contains the first FLR correction to the electromagnetic potential only:

$$H_2^{\text{dk}} = \alpha \frac{e^2}{2m_e c^2} A_{1\parallel}(\mathbf{X})^2.$$

2.2.1. Quasineutrality and Ampère equations

Both the quasineutrality and Ampère equations are derived using the functional derivative of the action with respect to ϕ_1 and $A_{1\parallel}$, i.e.

$$0 = \frac{\delta \mathcal{L}}{\delta \lambda} \circ \hat{\lambda}, \quad (8)$$

where λ and $\hat{\lambda}$ are respectively $\{\phi_1, A_{1\parallel}\}$ and $\{\hat{\phi}_1, \hat{A}_{1\parallel}\}$. Note that this is equivalent to solving the Euler-Lagrange equations. The weak form of the quasineutrality equation derived from the action equation (3) is then given by

$$\sum_{\sigma \neq e} \mathcal{Q}_{\sigma}^{\text{gyr}} + \mathcal{Q}_e^{\text{dk}} = \sum_{\sigma \neq e} \mathcal{Q}_{\sigma}^{\text{pol}}, \quad (9)$$

$$\mathcal{Q}_{\sigma}^{\text{gyr}} = \int d\Omega f_{\sigma} q_{\sigma} \langle \hat{\phi}_1 \rangle, \quad (10)$$

$$\mathcal{Q}_e^{\text{dk}} = - \int d\Omega f_e e \hat{\phi}_1(\mathbf{X}), \quad (11)$$

$$\mathcal{Q}_{\sigma}^{\text{pol}} = \epsilon_{\delta} \int d\Omega f_{\text{eq},\sigma} \frac{q_{\sigma}^2}{B} \frac{\partial}{\partial \mu} (\langle \phi_1 \hat{\phi}_1 \rangle - \langle \phi_1 \rangle \langle \hat{\phi}_1 \rangle), \quad (12)$$

where $\hat{\phi}_1$ represents an arbitrary test function, which in the case of ORB5 is a B-spline of a required order for the finite element discretization. On the left-hand side of the equation, $\mathcal{Q}_{\sigma}^{\text{gyr}}$ represents the gyro-charge density of the ions, $\mathcal{Q}_e^{\text{dk}}$ the drift-kinetic charge of the electrons and on the right-hand side, $\mathcal{Q}_{\sigma}^{\text{pol}}$ represents the

linear ion polarization charge. Note that due to the drift-kinetic approximation used for the electrons, there is no contribution to the polarization density from the electron species.

Similarly, the Ampère equation derived from the variational principle is given by

$$0 = -\epsilon_\delta \int \frac{dV}{4\pi} \nabla_\perp A_{1\parallel} \cdot \nabla_\perp \hat{A}_{1\parallel} + \sum_{\sigma \neq e} \int d\Omega f_\sigma \frac{q_\sigma p_z}{m_\sigma c} \langle \hat{A}_{1\parallel} \rangle \quad (13)$$

$$- \int d\Omega f_e \frac{e p_z}{m_e c} \hat{A}_{1\parallel}$$

$$- \epsilon_\delta \int d\Omega f_{eq,e} \left(\frac{e^2}{m_e c^2} A_{1\parallel} \hat{A}_{1\parallel} \right)$$

$$- \sum_{\sigma \neq e} \epsilon_\delta \int d\Omega f_{eq,\sigma} \left[\frac{q_\sigma^2}{m_\sigma c^2} A_{1\parallel} \hat{A}_{1\parallel} \right.$$

$$\left. + \frac{\mu}{2B} (A_{1\parallel} \nabla_\perp^2 \hat{A}_{1\parallel} + \hat{A}_{1\parallel} \nabla_\perp^2 A_{1\parallel}) \right],$$

for any test function $\hat{A}_{1\parallel}$. Note that in both the quasineutrality and Ampère equations, the polarization and magnetization terms come from the second order Hamiltonian H_2 .

2.2.2. Nonlinear gyrokinetic Vlasov equation

The gyrokinetic Vlasov equation for the distribution function f_σ of each species σ is reconstructed from the linearized gyrocenter characteristics according to the approximations performed on the action functional given by Eq. (3):

$$0 = \frac{df_\sigma}{dt} = \frac{\partial f_\sigma}{\partial t} + \dot{\mathbf{X}} \cdot \nabla f_\sigma + \dot{p}_z \frac{\partial f_\sigma}{\partial p_z}, \quad (14)$$

where the gyrocenter characteristics depend on the linearized Hamiltonian model:

$$\dot{\mathbf{X}} = \frac{c\hat{\mathbf{b}}}{q_\sigma B_\parallel^*} \times \nabla H + \frac{\partial H \mathbf{B}^*}{\partial p_z B_\parallel^*}, \quad (15)$$

$$\dot{p}_z = -\frac{\mathbf{B}^*}{B_\parallel^*} \cdot \nabla H, \quad (16)$$

with $H = H_0 + \epsilon_\delta H_1$, where H_0 is a Hamiltonian corresponding to the non perturbed guiding-center dynamics given by Eq. (4) and H_1 corresponds to the first order gyrocenter contributions given by Eq. (5).

For the ordering considered above, the characteristics become:

$$\dot{\mathbf{X}} = \frac{c\hat{\mathbf{b}}}{q_\sigma B_\parallel^*} \times \nabla \left[\mu B + \epsilon_\delta q_\sigma \left(\langle \phi_1 \rangle - \alpha \frac{p_z}{m_\sigma} \langle A_{1\parallel} \rangle \right) \right]$$

$$+ \frac{\mathbf{B}^*}{B_\parallel^*} \left(\frac{p_z}{m_\sigma} - \epsilon_\delta \alpha \frac{q_\sigma}{m_\sigma} \langle A_{1\parallel} \rangle \right), \quad (17)$$

$$\dot{p}_z = -\frac{\mathbf{B}^*}{B_\parallel^*} \cdot \nabla \left[\mu B + \epsilon_\delta q_\sigma \left(\langle \phi_1 \rangle - \alpha \frac{p_z}{m_\sigma} \langle A_{1\parallel} \rangle \right) \right], \quad (18)$$

which can be written in a different form to make the usual drift velocities appear:

$$\dot{\mathbf{X}} = \frac{p_z}{m_\sigma} \hat{\mathbf{b}} - \frac{c p_z^2}{q_\sigma m_\sigma B_\parallel^*} \left[\hat{\mathbf{b}} \times \left(\hat{\mathbf{b}} \times \frac{\nabla \times \mathbf{B}}{B} \right) \right] \quad (19)$$

$$+ \frac{c}{q_\sigma B_\parallel^*} \left(\mu B + \frac{p_z^2}{m_\sigma} \right) \hat{\mathbf{b}} \times \frac{\nabla B}{B}$$

$$+ \epsilon_\delta \frac{c}{B_\parallel^*} \hat{\mathbf{b}} \times \nabla \left(\langle \phi_1 \rangle - \alpha \frac{p_z}{m_\sigma} \langle A_{1\parallel} \rangle \right) + \epsilon_\delta \alpha \frac{c p_z}{m_\sigma B_\parallel^*} \langle A_{1\parallel} \rangle \boldsymbol{\kappa}$$

$$- \epsilon_\delta \alpha \frac{q_\sigma}{m_\sigma} \langle A_{1\parallel} \rangle \hat{\mathbf{b}}$$

$$\equiv \mathbf{v}_\parallel + \mathbf{v}_D + \mathbf{v}_{\nabla B} + \mathbf{v}_C + \mathbf{v}_{E \times B} + \mathbf{v}_{A_1}, \quad (20)$$

where $\boldsymbol{\kappa}$ is the curvature vector

$$\boldsymbol{\kappa} = \hat{\mathbf{b}} \times \left[\hat{\mathbf{b}} \times \frac{\nabla \times \mathbf{B}}{B} \right] + \frac{\nabla \mathbf{B} \times \hat{\mathbf{b}}}{B}. \quad (21)$$

The first term of the equation is the parallel velocity $\mathbf{v}_\parallel$, the second is the diamagnetic drift $\mathbf{v}_D$, the third term can be separated in the ∇B drift $\mathbf{v}_{\nabla B}$ and curvature drift $\mathbf{v}_C$, the fourth is the $E \times B$ drift $\mathbf{v}_{E \times B}$, and the last two terms are labeled as $\mathbf{v}_{A_1}$. Similarly, the same procedure can be applied to the p_z characteristic:

$$\dot{p}_z = \mu B \nabla \cdot \mathbf{B} - \frac{c p_z \mu}{q_\sigma B_\parallel^*} \left[\hat{\mathbf{b}} \times \left(\hat{\mathbf{b}} \times \frac{\nabla \times \mathbf{B}}{B} \right) \right] \cdot \nabla B \quad (22)$$

$$- \epsilon_\delta \nabla \left(\langle \phi_1 \rangle - \alpha \frac{p_z}{m_\sigma} \langle A_{1\parallel} \rangle \right) \cdot \left(q_\sigma \hat{\mathbf{b}} + \frac{c p_z}{B_\parallel^*} \boldsymbol{\kappa} \right) \quad (23)$$

$$\equiv -\frac{m_\sigma}{p_z} (\mathbf{v}_\parallel + \mathbf{v}_D + \mathbf{v}_C) \cdot \nabla \left(\mu B + \epsilon_\delta q_\sigma \langle \phi_1 \rangle - \epsilon_\delta \alpha \frac{p_z}{m_\sigma} \langle A_{1\parallel} \rangle \right). \quad (24)$$

In the ORB5 gyrokinetic model, different additional approximations can be made on the total time derivative operator introduced in Eq. (14): the linear and/or neoclassical limits. To this end, the characteristic equations (17) and (18) are slightly modified. In the linear limit, all the perturbed terms, proportional to ϵ_δ , are neglected leading to:

$$\dot{\mathbf{X}}^{\text{lin}} = \mathbf{v}_\parallel + \mathbf{v}_D + \mathbf{v}_{\nabla B} + \mathbf{v}_C, \quad (25)$$

$$\dot{p}_z^{\text{lin}} = -\mu \frac{m_\sigma}{p_z} (\mathbf{v}_\parallel + \mathbf{v}_D + \mathbf{v}_C) \cdot \nabla B. \quad (26)$$

2.3. Variants of the physical models

In this section, we present the different variants of the physical model presented above that are available in the ORB5 code. Usually, each variant can be obtained in the framework of the variational formulation by changing the H_0 , H_1 and H_2 Hamiltonians according to the corresponding approximations. This is the case for the long-wavelength approximated electromagnetic model as well as the electrostatic models with a Padé approximation and a strong background flow. For the adiabatic electron model, an external coupling of the gyrokinetic equations with a fluid polarization density of the electrons is assumed. Including this model into the general framework requires some additional approximations on the field term of the field-particles Lagrangian given by Eq. (3). Note that these models are not necessarily mutually exclusive and a summary of the different possible combinations will be presented at the end of the section.

2.3.1. Long wavelength approximation

This approximation is obtained by replacing the second order nonlinear Hamiltonian H_2 given by Eq. (7) in the gyrokinetic Lagrangian, Eq. (3), by the nonlinear Hamiltonian model [53] containing FLR expansions up to the second order for both its electrostatic and electromagnetic parts:

$$H_2^{\text{FLR}} = -\frac{m_\sigma c^2}{2B^2} |\nabla_\perp \phi_1(\mathbf{X})|^2 + \alpha \frac{q_\sigma^2}{2m_\sigma c^2} \left[A_{1\parallel}(\mathbf{X})^2 \right.$$

$$\left. + m_\sigma \left(\frac{c}{q_\sigma} \right)^2 \frac{\mu}{B} A_{1\parallel} \nabla_\perp^2 A_{1\parallel}(\mathbf{X}) \right]. \quad (27)$$

This changes only the term associated with the polarization charge of the quasineutrality equation, Eq. (9), so that Eq. (12) is replaced with

$$\mathcal{Q}_{\sigma, \text{LWA}}^{\text{pol}} = \epsilon_\delta \int d\Omega f_{eq,\sigma} \frac{m_\sigma c^2}{B^2} \nabla_\perp \phi_1 \cdot \nabla_\perp \hat{\phi}_1, \quad (28)$$

for any test functions $\widehat{\phi}_1$. The subscript LWA stands for long wavelength approximation. Since the magnetic terms in Eq. (27) remain unchanged compared to the Hamiltonian H_2 given by Eq. (7), for which the long wavelength approximation had already been considered, the corresponding Ampère equation remains the same as given by Eq. (13). The gyrokinetic Vlasov equation is unchanged as well, since the background H_0 and linear H_1 Hamiltonians are not affected by the approximation and no contributions from the second order Hamiltonian appear in the characteristics given by Eqs. (17) and (18).

2.3.2. Padé approximation

In addition to the long wavelength approximation, a Padé-approximated quasineutrality model for the ion species is available in ORB5 [30,54]. In practice, the Padé approximation is currently only implemented for one ion species ($\sigma = i$). In order to include this approximation inside the common variational principle, the linear Hamiltonian model has to be slightly modified with respect to Eq. (5) for both ions:

$$H_{1,\text{Padé}} = (1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}) H_1, \quad (29)$$

and electrons:

$$H_{1,\text{Padé}}^{\text{dk}} = (1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}) H_1^{\text{dk}}. \quad (30)$$

The nonlinear Hamiltonian model in that case is given by the FLR second-order truncated Hamiltonian H_2^{FLR} , Eq. (27). The quasineutrality equation in weak form is written in a different way by multiplying it by the operator $[1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}]$ to cancel the $[1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}]^{-1}$ term in the polarization density. This is done for computational reasons: the inverse of the block banded matrix coming from the discretization of the $[1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}]$ operator is a full matrix. For example, with drift-kinetic electrons, this leads to:

$$\begin{aligned} 0 &= q_i \int d\Omega f_i (1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}) (\widehat{\phi}_1) \\ &\quad - e \int d\Omega f_e (1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}) \widehat{\phi}_1 \\ &\quad + \epsilon_{\delta} \int d\Omega f_{\text{eq},i} \frac{m_i c^2}{B^2} \nabla_{\perp} \phi_1 \cdot \nabla_{\perp} \widehat{\phi}_1. \end{aligned} \quad (31)$$

2.3.3. Adiabatic electron model

In order to include a model with adiabatic electrons inside the variational formulation, we need to include a fluid approximation for the electron dynamics inside the field-particle Lagrangian. Compared to the main field-particle Lagrangian, Eq. (3), here the sum over the species in the first term is over the ion species only and the field term is modified by a purely electrostatic contribution from the electrons. The action principle for a model with adiabatic electrons is then given by

$$\begin{aligned} \mathcal{A}_{\text{adiab}} &= \int dt \mathcal{L}_{\text{adiab}} = \sum_{\sigma \neq e} \int dt d\Omega \left[\frac{q_{\sigma}}{c} \mathbf{A}^* \cdot \dot{\mathbf{X}} + \frac{m_{\sigma} c}{q_{\sigma}} \mu \dot{\phi} \right. \\ &\quad \left. - (H_0 + \epsilon_{\delta} H_1) \right] f_{\sigma} \\ &\quad + \epsilon_{\delta} \int dt dV \left[n_{e0} \phi_1 + \epsilon_{\delta} \frac{e}{2T_e} \right. \\ &\quad \left. \times n_{e0} (\phi_1 - \bar{\phi}_1)^2 \right] \\ &\quad - \epsilon_{\delta}^2 \sum_{\sigma \neq e} \int dt d\Omega H_2 f_{\text{eq},\sigma}, \end{aligned} \quad (32)$$

where $\bar{\phi}_1$ represents the flux-surface-averaged electric potential given by

$$\bar{\phi}_1 \equiv \frac{\int \phi_1 J(s, \theta^*) d\theta^* d\varphi}{\int J(s, \theta^*) d\theta^* d\varphi}, \quad (33)$$

where $J(s, \theta^*) = [\nabla s \cdot (\nabla \theta^* \times \nabla \varphi)]^{-1}$ is the Jacobian of the magnetic coordinate transformation and n_{e0} is the equilibrium electron density. Since the adiabatic electron model is only valid in the electrostatic limit, the velocity part of the phase space volume reduces to $dW = B_{\parallel}^* m_{\sigma} dv_{\parallel} d\mu$ and $\mathbf{B}^* = \nabla \times \mathbf{A}^*$ with $\mathbf{A}^* = \mathbf{A} + (c/q_{\sigma}) m_{\sigma} v_{\parallel} \hat{\mathbf{b}}$, while the spatial part $dV = d^3\mathbf{X}$ remains unchanged with respect to the electromagnetic case. The Hamiltonian models are now defined for a simplified electrostatic case as

$$H_0 = \frac{m_{\sigma} v_{\parallel}^2}{2} + \mu B, \quad (34)$$

$$H_1 = q_{\sigma} \langle \phi_1 \rangle. \quad (35)$$

The nonlinear ion dynamics is defined by the electrostatic part of either the full FLR, the Padé-approximated, or the second order FLR long-wavelength-approximated nonlinear Hamiltonian.

The corresponding Vlasov equation does not contain any contribution from the electron species, so we have for ions ($\sigma = i$)

$$0 = \frac{df_{\sigma}}{dt} = \frac{\partial f_{\sigma}}{\partial t} + \dot{\mathbf{X}} \cdot \nabla f_{\sigma} + \dot{v}_{\parallel} \frac{\partial f_{\sigma}}{\partial v_{\parallel}}, \quad (36)$$

with the characteristics corresponding to the electrostatic limit ($\alpha = 0$) of Eqs. (15) and (16):

$$\dot{\mathbf{X}} = \frac{c\hat{\mathbf{b}}}{q_{\sigma} B_{\parallel}^*} \times \nabla (\mu B + \epsilon_{\delta} q_{\sigma} \langle \phi_1 \rangle) + \frac{\mathbf{B}^*}{B_{\parallel}^*}, \quad (37)$$

$$\dot{v}_{\parallel} = -\frac{\mathbf{B}^*}{B_{\parallel}^*} \cdot \nabla (\mu B + \epsilon_{\delta} q_{\sigma} \langle \phi_1 \rangle).$$

For the quasineutrality equation, only the electron contribution to the gyro-charge term is modified, leading to

$$Q_{e,\text{adiab}}^{\text{dk}} = \epsilon_{\delta} \int dV \frac{en_{e0}}{T_e} (\phi_1 - \bar{\phi}_1) \widehat{\phi}_1 + \int dV n_{e0} \widehat{\phi}_1. \quad (38)$$

2.3.4. Hybrid electron model

There is also the possibility to include a hybrid electron model inside the variational formulation. In that case the fraction of passing electrons designated with a coefficient α_p is treated as an adiabatic species, while the fraction of trapped electrons is treated as a drift-kinetic species. At the same time, the ions are treated as kinetic species. The corresponding action functional is given by

$$\begin{aligned} \mathcal{A}_{\text{hybrid}} &= \int dt \mathcal{L}_{\text{hybrid}} = \sum_{\sigma \neq e} \int dt d\Omega \left[\frac{q_{\sigma}}{c} \mathbf{A}^* \cdot \dot{\mathbf{X}} + \frac{m_{\sigma} c}{q_{\sigma}} \mu \dot{\phi} \right. \\ &\quad \left. - (H_0 + \epsilon_{\delta} H_1) \right] f_{\sigma} \\ &\quad - \epsilon_{\delta}^2 \sum_{\sigma \neq e} \int dt d\Omega H_2 f_{\text{eq},\sigma} \\ &\quad + \int dt dV \int_{\text{trapped}} dW \left[\frac{e}{c} \mathbf{A}^* \cdot \dot{\mathbf{X}} \right. \\ &\quad \left. + \frac{m_e c}{e} \mu \dot{\phi} - (H_0 + \epsilon_{\delta} H_1^{\text{dk}}) \right] f_e \\ &\quad + \alpha_p \epsilon_{\delta} \int dt dV \left[n_{e0} \phi_1 + \epsilon_{\delta} \frac{e}{2T_e} \right. \\ &\quad \left. \times n_{e0} (\phi_1 - \bar{\phi}_1)^2 \right], \end{aligned} \quad (39)$$

where the integral over the fraction of trapped electrons in the velocity phase space is assumed with $\int_{\text{trapped}} dW$. The phase space configuration is the same as in the case of an adiabatic electron model. The gyrocenter model used for modeling the ion species dynamics is identical to the one presented for the adiabatic electron model discussed in the previous section, i.e. the Hamiltonians H_0 and H_1 are given by Eqs. (34)–(35) and the nonlinear Hamiltonian is coming from either the full FLR, the Padé approximation or the long wavelength approximation. Concerning the gyrocenter models used for modeling the hybrid electron dynamics, the equilibrium dynamics is defined with H_0 given by Eq. (34). The linear part of the trapped electron dynamics is defined by the drift-kinetic model defined by Eq. (6) with $\alpha = 0$. The quasineutrality equation is only affected through the electron contribution to the gyro-charge term that reads

$$Q_{e,\text{hyb}}^{\text{dk}} = \alpha_p \epsilon_\delta \int dV n_{e0} \frac{e}{T_e} (\phi_1 - \bar{\phi}_1) \hat{\phi}_1 + \int dV n_{e0} \hat{\phi}_1 + \int_{\text{trapped}} d\Omega f_e \hat{\phi}_1. \quad (40)$$

The ion characteristics are reconstructed identically to the case with adiabatic electrons, accordingly to Eq. (37). The characteristics for the electrons are defined by the simplified drift-kinetic equations corresponding to the dynamics of $H = H_0 + H_1^{\text{dk}}$:

$$\dot{\mathbf{X}} = \frac{\hat{\mathbf{c}}\mathbf{b}}{q_\sigma B_\parallel^*} \times \nabla (\mu B + \epsilon_\delta e \phi_1) + \frac{\mathbf{B}^*}{B_\parallel^*} v_\parallel, \quad (41)$$

$$\dot{v}_\parallel = -\frac{\mathbf{B}^*}{B_\parallel^*} \cdot \nabla (\mu B + \epsilon_\delta e \phi_1).$$

Note that all the electrons, passing and trapped, are evolved according to the above drift-kinetic characteristics.

The hybrid electron model presented above was originally implemented to simulate linear electron modes such as TEM, allowing for a larger timestep than with fully drift-kinetic electrons. However, in nonlinear regime, it does not ensure the ambipolar condition – which also impacts the conservation of the toroidal angular momentum – as no flux-surface-averaged passing-electron density is accounted for. Spurious sources of e.g. density and momentum are added in this model because particles that are becoming trapped (detrapped) are suddenly (not) accounted for kinetically in the quasi-neutrality equation. The trapping/detrapping processes are the collisions and parallel nonlinearity that affect the particle pitch angle and radial drifts that change the value of the local background magnetic field. To address this issue, an upgraded hybrid electron model has been implemented in ORB5 as an improvement of the model presented in [55]. The idea of this updated model is to take into consideration only the $n = m = 0$ component of the passing electron density while keeping an adiabatic response for the other passing components. This way, the quasineutrality equation, Eq. (40), is slightly changed as follows

$$Q_{e,\text{hyb}}^{\text{dk}} = \alpha_p \epsilon_\delta \int dV n_{e0} \frac{e}{T_e} (\phi_1 - \bar{\phi}_1) \hat{\phi}_1 + \int dV n_{e0} \hat{\phi}_1 + \int_{\text{trapped}} d\Omega f_e \hat{\phi}_1 + \int_{\text{passing}} d\Omega f_e \hat{\phi}_1^{00}, \quad (42)$$

where $\hat{\phi}_1^{00}$ is the $n = m = 0$ component of the arbitrary test function $\hat{\phi}_1$.

2.3.5. Summary of the models

All the variants of the particle models presented in the previous sections are summarized here. The changes brought by

the different models mainly come through the quasineutrality equation which can be written

$$\sum_{\sigma \neq e} Q_\sigma^{\text{gyr}} + Q_e^{\text{dk}} = \sum_{\sigma \neq e} Q_\sigma^{\text{pol}}, \quad (43)$$

$$Q_\sigma^{\text{gyr}} = \int d\Omega f_\sigma q_\sigma \langle \hat{\phi}_1 \rangle, \quad (44)$$

$$Q_e^{\text{dk}} = - \int d\Omega f_e e \hat{\phi}_1(\mathbf{X}), \quad (45)$$

$$Q_\sigma^{\text{pol}} = \epsilon_\delta \int d\Omega f_{\text{eq},\sigma} \frac{q_\sigma^2}{B} \frac{\partial}{\partial \mu} (\langle \phi_1 \hat{\phi}_1 \rangle - \langle \phi_1 \rangle \langle \hat{\phi}_1 \rangle), \quad (46)$$

where Q_σ^{gyr} is the term corresponding to the ion gyrodensity contribution, Q_e^{dk} is the term corresponding to the electron drift kinetic density contribution and Q_σ^{pol} represents the polarization density contribution from the ions. For the ions, only Q_σ^{pol} is affected by the different models:

$$Q_{\sigma,\text{LWA}}^{\text{pol}} = \epsilon_\delta \int d\Omega f_{\text{eq},\sigma} \frac{m_\sigma c^2}{B^2} \nabla_\perp \phi_1 \cdot \nabla_\perp \hat{\phi}_1, \quad (47)$$

$$Q_{i,\text{Padé}}^{\text{pol}} = \epsilon_\delta \int d\Omega f_{\text{eq},i} \frac{m_i c^2}{B^2} \left[1 - \nabla_\perp \cdot \rho_i^2 \nabla_\perp \right]^{-1} \left[\nabla_\perp \phi_1 \cdot \nabla_\perp \hat{\phi}_1 \right]. \quad (48)$$

Note that in the case of the Padé approximation, all the quasineutrality equation is multiplied by $\left[1 - \nabla_\perp \cdot \rho_i^2 \nabla_\perp \right]$ to avoid having to invert it.

For the electrons, only Q_e^{dk} is changed by the different fluid and hybrid approximations:

$$Q_{e,\text{adiab}}^{\text{dk}} = \epsilon_\delta \int dV \frac{en_{e0}}{T_e} (\phi_1 - \bar{\phi}_1) \hat{\phi}_1 - \int dV n_{e0} \hat{\phi}_1, \quad (49)$$

$$Q_{e,\text{hyb}}^{\text{dk}} = \alpha_p \epsilon_\delta \int dV \frac{en_{e0}}{T_e} (\phi_1 - \bar{\phi}_1) \hat{\phi}_1 - \int dV n_{e0} \hat{\phi}_1 - e \int_{\text{trapped}} d\Omega f_e \hat{\phi}_1 + \int_{\text{passing}} d\Omega f_e \hat{\phi}_1^{00}. \quad (50)$$

In ORB5, the previous approximations are not mutually exclusive, i.e. each model for the polarization density can be combined with any electron model.

2.4. δf method and background distribution functions

The ORB5 code uses a δf control-variates approach to reduce the numerical noise due to the finite phase-space sampling [36,56]. The rationale of this method is to separate the total distribution function into two parts: a time-independent part f_0 and a time-dependent part δf . The first function, f_0 , is supposed to be known and easily computable. Only the δf part is represented with a sample of “numerical particles” or “markers”. The statistical sampling error will thus be reduced, as compared to a full- f method, if $|\delta f| \ll f_0$.

In the collisionless limit and in the absence of sources, the total distribution function is conserved along the trajectories. Using the δf separation, we obtain

$$\frac{d\delta f}{dt} = -\frac{df_0}{dt} = -\frac{df_0}{dt} \Big|_0 - \frac{df_0}{dt} \Big|_1, \quad (51)$$

where the time-derivative operator has been split into two parts labeled by 0 and 1 and which respectively represent the unperturbed dynamics, i.e. without the fluctuating fields, and the perturbed dynamics. In the *standard* δf method, we choose $f_0 \equiv f_{\text{eq}}$ to be an equilibrium distribution, solution of the unperturbed collisionless equations of motion and thus, to satisfy $\{f_{\text{eq}}, H_0\} = 0$ reducing Eq. (51) to

$$\frac{d\delta f}{dt} = -\frac{df_{\text{eq}}}{dt} \Big|_1. \quad (52)$$

In ORB5, different choices for the initial distribution function are available. The plasma can be assumed to be in a local thermodynamic equilibrium described by a *local Maxwellian* $f_L(\psi, \epsilon, \mu)$. Both the particle energy ϵ and the magnetic moment μ are constants of motion but the poloidal magnetic flux ψ is not. The local Maxwellian is therefore not invariant under the unperturbed dynamics and Eq. (51) must be used. The inclusion of the $df_L/dt|_0$ term leads to the drive of a spurious zonal flow discussed in [10], which appears already in the linear phase of a simulation, even though zonal flows are linearly stable and excited through nonlinear coupling [11]. As done in many PIC codes, the above-mentioned term responsible for this zonal flow drive can be neglected but it is not consistent with the perturbative ordering used here.

The alternative approach is to use a distribution function that is a true equilibrium, implying that f_0 is a function of constants of motion only. This verified by the so-called *canonical Maxwellian* $f_C(\psi_0, \epsilon, \mu)$, where $\psi_0 = \psi + (m_\sigma c/q_\sigma)(F(\psi)/B)v_\parallel$ is the toroidal momentum which is conserved in an axisymmetric toroidal system. However, it is easily shown that the effective density and temperature computed from f_C are different from the ones given as input and function of ψ_0 . The use of a canonical Maxwellian can lead to large, unrealistic values of parallel flows preventing any instability to develop, especially for small system size and large n_0 and $v_\parallel$ gradients [57]. To address this issue, a *corrected canonical Maxwellian* $f_{CC}(\hat{\psi}, \epsilon, \mu)$ is used. A correction term is added to the toroidal momentum to minimize the gap between the local and canonical Maxwellians while still being an equilibrium distribution. The corrected toroidal momentum reads

$$\hat{\psi} = \psi_0 + \psi_{0,\text{corr}} = \psi_0 - \text{sign}(v_\parallel) \frac{m_\sigma c}{q_\sigma} R_0 \sqrt{2(\epsilon - \mu B_0)} \mathcal{H}(\epsilon - \mu B_0), \quad (53)$$

where R_0 is the major radius and $\mathcal{H}$ is the Heaviside function. The correction term is zero for trapped particles and of opposite sign for forward and backward passing particles. The corrected toroidal momentum being built only with constants of motion, $f_{CC}(\hat{\psi}, \epsilon, \mu)$ satisfies $\{f_{CC}, H_0\} = 0$.

2.5. Strong flows

The strong flow gyrokinetic ordering allows for $u_E/v_{\text{th},i} \sim 1$, with $\mathbf{u}_E = c(\hat{\mathbf{b}} \times \nabla \Phi/B)$ the background $E \times B$ velocity, where Φ represents the background electric potential, and $v_{\text{th},i} = \sqrt{T_i/m_i}$ is the ion thermal velocity [58]. Implementing this ordering in ORB5 enables the treatment of plasmas rotating toroidally at close to the Mach velocity. More details of this formalism have been published earlier [33]. In this case, a further approximation is performed on the background distribution function. While a local Maxwellian is used for the polarization density in the quasineutrality equation, the canonical Maxwellian is implemented for the derivation of the gyrokinetic Vlasov equation.

In order to include the model containing a background electrostatic potential Φ within the general field-gyrocenter action given by Eq. (3), the background Hamiltonian H_0 , as well as the symplectic magnetic potential $\mathbf{A}^*$ have to be consistently modified:

$$H_0^{\text{flow}} = q_\sigma \Phi + \mu B + \frac{p_z^2 + (m_\sigma \mathbf{u}_E)^2}{2m_\sigma}, \quad (54)$$

$$\text{and } \mathbf{A}^* = \mathbf{A} + (c/q_\sigma) p_z \hat{\mathbf{b}} + (m_\sigma c/q_\sigma) \mathbf{u}_E.$$

In ORB5, when considering strong flows, only the electrostatic limit is considered. This corresponds to setting $\alpha = 0$ in Eq. (3). Remark that including the background $E \times B$ velocity does not affect the quasineutrality equation, since no corrections due to

the presence of a strong flow are included into the linear and nonlinear Hamiltonian models given by Eqs. (5)–(7). The gyrokinetic Vlasov equation is modified according to the change in the background dynamics from H_0 given by Eq. (4) to H_0^{flow} given by Eq. (54). The corresponding δf gyrokinetic Vlasov equation is reconstructed from the modified characteristics. Since the perturbed magnetic field is not considered, $p_z = m_\sigma v_\parallel$ is a purely kinetic momentum:

$$\begin{aligned} \dot{\mathbf{X}} &= \frac{c\hat{\mathbf{b}}}{q_\sigma B_\parallel^*} \times \nabla \left(q_\sigma \Phi + \mu B + \frac{m_\sigma}{2} |\mathbf{u}_E|^2 + \epsilon_\delta q_\sigma \langle \phi_1 \rangle \right) \\ &\quad + \frac{\mathbf{B}^*}{B_\parallel^*} v_\parallel, \\ \dot{v}_\parallel &= -\frac{\mathbf{B}^*}{B_\parallel^*} \cdot \nabla \left(q_\sigma \Phi + \mu B + \frac{m_\sigma}{2} |\mathbf{u}_E|^2 + \epsilon_\delta q_\sigma \langle \phi_1 \rangle \right). \end{aligned} \quad (55)$$

For strongly rotating plasmas, with Mach number around one, dynamic pressure resulting from the flow is comparable to the thermal pressure, and a modified Grad–Shafranov equation should be used to accurately compute the magnetic equilibrium. To self-consistently include these effects, we have used the MHD code FLOW [59] which can solve the MHD force balance equation in the presence of a background flow. FLOW reads the equilibrium via the standard EQDSK format [50]. We have considered only toroidally rotating MHD equilibria, with the temperature being a flux surface function, as this allows collisionless kinetic and MHD equilibria to be consistent in the large-system size limit.

2.5.1. Global gyrokinetic equilibria for rotating plasmas

The constants of motion are the magnetic moment μ , the unperturbed energy of the particle $\epsilon = H_0^{\text{flow}}$, the sign of the parallel velocity (for passing particles), and finally the toroidal canonical momentum ψ_0 , which is conserved in an tokamak due to axisymmetry. The strong-flow canonical momentum ψ_C^{flow} is an extension of the canonical momentum discussed in Section 2.4:

$$\psi_C^{\text{flow}} = \psi + \frac{m_\sigma c}{q_\sigma} \frac{F}{B} v_\parallel + \frac{m_\sigma c}{q_\sigma} u_\varphi, \quad (56)$$

where u_φ is the toroidal component of the background $E \times B$ velocity.

Background distribution functions appropriate for pure poloidal rotation and pure toroidal rotation are implemented, but only the most common case of a toroidally rotating plasma is described here, and the reader is referred to Ref. [60] for more details. In the presence of toroidal rotation, the canonical Maxwellian, which is corrected so that the flux surface averaged density remains close to n_0 when rotation is introduced, is given by

$$\begin{aligned} f_C &= \left(\frac{m_\sigma}{2\pi T_0(\psi_C^{\text{flow}})} \right)^{3/2} n_0(\psi_C^{\text{flow}}) \exp \left\{ -\frac{1}{T_0(\psi_C^{\text{flow}})} \right. \\ &\quad \times \left[H_0^{\text{flow}} + \frac{m_\sigma R_0 (\psi_C^{\text{flow}})^2}{2} \left(\frac{\partial \bar{\Phi}}{\partial \psi} \right)^2 \right] \Bigg\}, \end{aligned} \quad (57)$$

where $\bar{\Phi}$ is the flux surface average of Φ . In the local limit, this choice leads to an in-out density variation

$$n_\sigma = n_0(\psi) \exp \left[\frac{m_\sigma (R^2 - R_0^2) \Omega^2}{2} \right], \quad (58)$$

where the plasma rotation frequency Ω may be expressed as $\Omega = \partial \Phi / \partial \psi$.

2.6. Collisions

The inclusion of collisions in a gyrokinetic code like ORB5 is important to assess the right level of transport. Indeed, collisions are required to model the neoclassical physics, which is a key player in the transport of certain classes of particles, e.g. heavy impurities. Furthermore, collisions are known to impact turbulence. For example, ITG driven turbulence increases when collisions are taken into account due to the collisional damping of the zonal flows [38,61]. On the other hand, TEM turbulence is reduced by collisions via the collisional detrapping of electrons.

ORB5 currently includes ion–ion intra-species and electron–ion collisions [27]. For the collisional dynamics, FLR and polarization effects are neglected. In ORB5, collisions are represented by a linearized Landau collision operator. The linearization procedure is done with respect to a local Maxwellian background f_L . The full Landau operator describing collisions of species a on species b is bilinear with respect to the distribution functions f_a and f_b and may thus be decomposed into four terms: $C_{ab}[f_b, f_a] = C_{ab}[f_{b,L}, f_{a,L}] + C_{ab}[f_{b,L}, \delta f_a] + C_{ab}[\delta f_b, f_{a,L}] + C_{ab}[\delta f_b, \delta f_a]$, where δf_σ is the perturbed part of the distribution of species σ . Note that in our notation, $C_{ab}[f_a, f_b]$ refers to the collisions of species a on species b . For the whole collision part, the species background distribution function is converted to a local Maxwellian if it is not already the case. After, the collision dynamics has been treated the background Maxwellian is converted back to its original form if needed, $f = f_0 + \delta f = f_{0L} + \delta f_L$. The perturbed part δf is modified accordingly such that the total distribution remains unchanged in this “background switching” process. For Maxwellian distributions with identical parallel velocities and temperatures, the first term on the right-hand side is zero. Assuming the perturbation is small, the final, nonlinear term, is also neglected leaving two terms called the “test particle” term $C_{TP}[f_{b,L}, \delta f_a]$ and the “background reaction” term $C_{BR}[\delta f_b, f_{a,L}]$.

For the self-collisions, the “test particle” term can be readily evaluated using the exact Landau operator in its drag-diffusion form:

$$C_{TP}[f_L, \delta f] = \frac{\partial}{\partial \mathbf{v}} \cdot [\mathbf{\Gamma}(f_L) \delta f] - \frac{\partial^2}{\partial \mathbf{v} \mathbf{v}} : [\bar{\bar{D}}(f_L) \delta f], \quad (59)$$

where the drag vector and the diffusion tensor are respectively given by

$$\mathbf{\Gamma} = -\bar{v} H(x) \mathbf{v}, \quad \bar{\bar{D}} = \frac{\bar{v} v_{th}^2}{4} \left[K(x) \left(\bar{\bar{I}} - \frac{\mathbf{v} \cdot \mathbf{v}}{v^2} \right) + 2H(x) \frac{\mathbf{v} \cdot \mathbf{v}}{v^2} \right], \quad (60)$$

where the collision frequency is defined as $\bar{v} = 8\pi n q^4 \ln \Lambda / m^2 v_{th}^3$, $x = v / \sqrt{2} v_{th}$ is the normalized velocity with $v_{th} = \sqrt{T/m}$ the thermal velocity of the species, and $\bar{\bar{I}}$ is the identity tensor. The Coulomb logarithm $\ln \Lambda$, is assumed constant across the plasma, and typically has a value of 10–15. The functions $K(x)$ and $H(x)$ are resulting from the analytical evaluation of the Rosenbluth potentials in the case of a Maxwellian background distribution:

$$H(x) = \frac{1}{2\sqrt{2}x^3} [\text{erf}(x) - x \text{erf}'(x)], \quad (61)$$

$$K(x) = \frac{1}{\sqrt{2}x} \phi(x) - H(x). \quad (62)$$

where erf stands for the error function.

Evaluating the background reaction term exactly would in particular require the reconstruction of the δf distribution function and taking its velocity derivative. Such a direct approach is too expensive and includes steps subject to significant noise in a

PIC code. Instead, ORB5 uses an approximation first suggested by [62]: $C_{BR}[\delta f, f_L] \simeq f_L \beta(\delta f)$, with

$$\beta(\delta f) = \frac{1}{n} \left[6\sqrt{\pi} H(x) \frac{\delta P_{\parallel v}}{v_{th}^2} + \sqrt{\pi} G(x) \frac{\delta E}{v_{th}^2} \right], \quad (63)$$

where $G(v) = (4x^2 - 1)H(x) - K(x)$. The two terms $\delta P_{\parallel}$ and δE represent respectively the parallel momentum and energy transferred to the distribution by the “test particle” operator:

$$\delta P_{\parallel} = - \int C_{TP}[f_L, \delta f] v_{\parallel} d^3 v, \quad (64)$$

$$\delta E = - \int C_{TP}[f_L, \delta f] v^2 d^3 v. \quad (65)$$

This approximation can be shown to satisfy the desirable properties of a collision operator [62,63]. Indeed, it conserves the mass and, when combined with its counterpart $C_{TP}[f_L, \delta f]$, conserves also the momentum and energy. Furthermore, the combined linear operator is self-adjoint and satisfies the H-theorem. The operator is zero if the perturbation is a shifted linearized Maxwellian, i.e. such distributions are stationary states.

The only interspecies collisions which are currently taken into account in ORB5 are the electron–ion collisions. The “test particle” part of the electron–ion collisions in ORB5 is represented by a Lorentz operator, which assumes a large mass ratio between ions and electrons. In this limit, electrons experience only pitch-angle scattering. This Lorentz operator can simply be written:

$$C_{ei}[f_i, \delta f_e] = -v_{ei}(v) \frac{\partial}{\partial \xi} \left[(1 - \xi^2) \frac{\partial \delta f_e}{\partial \xi} \right], \quad (66)$$

where the electron–ion collision frequency is given by $v_{ei}(v) = (\bar{v}_{ei}/4)(v_{th,e}/v)^3$, with $\bar{v}_{ei} = 8\pi n_i Z^2 e^4 \ln \Lambda / m_e^2 v_{th,e}^3$ and $\xi = v_{\parallel}/v$ is the pitch angle variable. The “test particle” Lorentz operator conserves the mass and energy. The “background reaction” of the Lorentz operator is neglected in ORB5. Therefore, momentum conservation is not ensured by the reduced electron–ion collision operator.

In the local neoclassical limit, which is made neglecting the fluctuating electromagnetic fields and assuming small banana widths as compared to the characteristic lengths of the system, the Fokker–Planck equation reduces to

$$\left. \frac{d\delta f}{dt} \right|_{\text{neo}} + C_{TP}[\delta f] = -\frac{d f_{LM}}{dt} - C_{BR}[f_{LM}], \quad (67)$$

where the first term on the left hand side is defined using the δf characteristics in the neoclassical limit, i.e. neglecting all drift velocities compared to the parallel drift velocity:

$$\dot{\mathbf{x}}^{\text{neo}} = \mathbf{v}_{\parallel}, \quad (68)$$

$$\dot{p}_z^{\text{neo}} = -\mu \frac{m_{\sigma}}{p_z} \mathbf{v}_{\parallel} \cdot \nabla B. \quad (69)$$

On the other hand, the first term on the right hand side, describing the background dynamics, contains all the drifts, Eqs. (17) and (18).

2.7. Conservation laws and diagnostics

In this section we present the conserved quantities associated with the field-particle Lagrangian, which are implemented in ORB5 as diagnostic tools. These quantities can be obtained from a direct application of the Noether method, details of the derivation can be found in [52]. We start by presenting the energy invariant corresponding to each model. This invariant is used for constructing the so-called power balance diagnostic, which allows one to verify the quality of numerical simulations.

First, we provide a generic expression for the energy density corresponding to the most complete electromagnetic model,

which can also be obtained from a direct application of the Noether method, see e.g. [52].

$$\begin{aligned} \mathcal{E}^{\text{EM}} = & \sum_{\sigma} \int d\Omega H_0 f_{\sigma} + \epsilon_{\delta} \sum_{\sigma \neq e} \int d\Omega H_1 f_{\sigma} \\ & + \epsilon_{\delta} \int d\Omega H_1^{\text{dk}} f_e \\ & + \epsilon_{\delta}^2 \sum_{\sigma \neq e} \int d\Omega H_2 f_{\text{eq},\sigma} + \alpha \epsilon_{\delta}^2 \int d\Omega H_2^{\text{dk}} f_{\text{eq},e} \\ & + \alpha \epsilon_{\delta}^2 \int dV \frac{|\nabla_{\perp} A_{1\parallel}|^2}{8\pi}. \end{aligned} \quad (70)$$

This expression can be simplified and rewritten in the form of code diagnostics by direct substitution of the expression for the Hamiltonians H_0 , H_1 given by Eqs. (4) and (5), while H_2 is given by Eq. (7) in the case of the all-orders in FLR polarization density model and by Eq. (27) in the case of the long-wavelength approximation. All the terms in the expression of the energy except the first one are then rewritten using the corresponding quasineutrality and Ampère equations in their weak form. Here we choose a particular test function $\hat{\phi}_1 = \phi_1$ and we substitute it in Eqs. (9)–(12) or, for the case of the long-wavelength approximation, in Eqs. (9), (10), (11), and (28). Similarly, the test function $\hat{A}_{1\parallel} = A_{1\parallel}$ is substituted to the corresponding Ampère equation given by Eq. (13). In PIC codes, particles and fields are evaluated in two different ways: particles are advanced continuously along their characteristics while fields are evaluated with finite elements on a fixed grid. To test the quality of the simulation, the contributions to the energy from the particles and from the fields should be computed independently. This is why we are considering the power balance equation, also called the $E \times B$ transfer equation. The code diagnostics are implemented to verify the following balance equation for $\mathcal{E}^{\text{EM}} = \mathcal{E}_{\text{F}} + \mathcal{E}_{\text{kin}}$:

$$0 = \frac{d\mathcal{E}^{\text{EM}}}{dt} \Rightarrow \frac{d\mathcal{E}_{\text{kin}}}{dt} = -\frac{d\mathcal{E}_{\text{F}}}{dt}, \quad (71)$$

where the time derivative of the left hand side of this relation can be evaluated through the particle characteristics and the right hand side from the field contributions evaluated on the grid.

From Eq. (70), the first term on the r.h.s. is defined as the “kinetic energy” $\mathcal{E}_{\text{kin}}$:

$$\mathcal{E}_{\text{kin}} = \sum_{\sigma} \int d\Omega H_0 f_{\sigma} = \sum_{\sigma} \int d\Omega \left(\frac{p_z^2}{2m_{\sigma}} + \mu B \right) f_{\sigma}, \quad (72)$$

which depends only on the unperturbed Hamiltonian H_0 and therefore, its time derivative can be evaluated considering only the unperturbed characteristics. The other terms are defined as the “field energy” $\mathcal{E}_{\text{F}}$, which can be written, for the case of the H_2 Hamiltonian written in the LWA, Eq. (27), as:

$$\begin{aligned} \mathcal{E}_{\text{F}} = & \epsilon_{\delta} \sum_{\sigma \neq e} \int d\Omega q_{\sigma} \left\langle \phi_1 - \alpha A_{1\parallel} \frac{p_z}{m_{\sigma}} \right\rangle f_{\sigma} \\ & - \epsilon_{\delta} \int d\Omega e \left(\phi_1 - \alpha A_{1\parallel} \frac{p_z}{m_e} \right) f_e \\ & + \epsilon_{\delta}^2 \sum_{\sigma \neq e} \int d\Omega f_{\text{eq},\sigma} \left\{ -\frac{m_{\sigma} c^2}{2B^2} |\nabla_{\perp} \phi_1|^2 + \alpha \frac{q_{\sigma}^2}{2m_{\sigma}} \right. \\ & \times \left[A_{1\parallel}^2 + \left(\frac{m_{\sigma}}{q_{\sigma}} \right)^2 \frac{\mu}{B} A_{1\parallel} \nabla_{\perp}^2 A_{1\parallel} \right] \Big\} \\ & + \epsilon_{\delta}^2 \alpha \int d\Omega f_{\text{eq},e} \frac{e^2}{2m_e} A_{1\parallel}^2 + \epsilon_{\delta}^2 \alpha \int dV \frac{|\nabla_{\perp} A_{1\parallel}|^2}{8\pi}. \end{aligned} \quad (73)$$

$$\begin{aligned} & \times \left[A_{1\parallel}^2 + \left(\frac{m_{\sigma}}{q_{\sigma}} \right)^2 \frac{\mu}{B} A_{1\parallel} \nabla_{\perp}^2 A_{1\parallel} \right] \Big\} \\ & + \epsilon_{\delta}^2 \alpha \int d\Omega f_{\text{eq},e} \frac{e^2}{2m_e} A_{1\parallel}^2 + \epsilon_{\delta}^2 \alpha \int dV \frac{|\nabla_{\perp} A_{1\parallel}|^2}{8\pi}. \end{aligned} \quad (74)$$

$$+ \epsilon_{\delta}^2 \alpha \int d\Omega f_{\text{eq},e} \frac{e^2}{2m_e} A_{1\parallel}^2 + \epsilon_{\delta}^2 \alpha \int dV \frac{|\nabla_{\perp} A_{1\parallel}|^2}{8\pi}. \quad (75)$$

Using the quasineutrality equations (9)–(11) with the polarization term in the LWA, Eq. (28), Ampère equation, Eq. (13), and setting $\hat{\phi}_1 = \phi_1$ and $\hat{A}_{1\parallel} = A_{1\parallel}$, we obtain two equivalent expressions for the field energy:

$$\begin{aligned} \mathcal{E}_{\text{F}} = & \epsilon_{\delta} \frac{1}{2} \sum_{\sigma \neq e} q_{\sigma} \int d\Omega \left(\langle \phi_1 \rangle - \alpha \frac{p_z}{m_{\sigma}} \langle A_{1\parallel} \rangle \right) f_{\sigma} \\ & - \epsilon_{\delta} \frac{1}{2} e \int d\Omega \left(\phi_1 - \alpha \frac{p_z}{m_e} A_{1\parallel} \right) f_e. \end{aligned} \quad (76)$$

Note that Eq. (76) does not depend on the particular choice for the nonlinear Hamiltonian H_2 . Indeed, Eq. (76) is also valid for the all order FLR polarization density, Eq. (12). This is a direct consequence of the fact that the equations of motion, which are used for rewriting the expression of the energy are obtained from the same field-particle Lagrangian.

Similarly, a second expression for the field energy written in terms of the polarizations and magnetizations and depending on the expression of the nonlinear Hamiltonian H_2 can be obtained. For the full FLR polarization density given by Eq. (12), the alternative field energy is given by

$$\begin{aligned} \mathcal{E}_{\text{F}} = & \frac{1}{2} \sum_{\sigma \neq e} \epsilon_{\delta} \int d\Omega f_{\text{eq},\sigma} \frac{q_{\sigma}^2}{B} \frac{\partial}{\partial \mu} \left\langle \tilde{\phi}_1 (\mathbf{X} + \boldsymbol{\rho}_0)^2 \right\rangle \\ & + \frac{1}{2} \alpha \sum_{\sigma \neq e} \epsilon_{\delta}^2 \int d\Omega f_{\text{eq},\sigma} \left(\frac{q_{\sigma}^2}{m_{\sigma}} A_{1\parallel}^2 + \frac{\mu}{B} A_{1\parallel} \nabla_{\perp}^2 A_{1\parallel} \right) \end{aligned} \quad (77)$$

$$+ \frac{1}{2} \alpha \epsilon_{\delta}^2 \int d\Omega f_{\text{eq},e} \frac{e^2}{m_e} A_{1\parallel}^2 + \alpha \epsilon_{\delta}^2 \int dV \frac{dV}{8\pi} |\nabla_{\perp} A_{1\parallel}|^2. \quad (78)$$

For the polarization density in the LWA, Eq. (28), the field energy becomes

$$\begin{aligned} \mathcal{E}_{\text{F}}^{\text{LWA}} = & \frac{1}{2} \sum_{\sigma \neq e} \epsilon_{\delta} \int d\Omega \frac{m_{\sigma} c^2}{B^2} f_{\text{eq},\sigma} |\nabla_{\perp} \phi_1|^2 \\ & + \frac{1}{2} \alpha \sum_{\sigma \neq e} \epsilon_{\delta}^2 \int d\Omega f_{\text{eq},\sigma} \left(\frac{q_{\sigma}^2}{m_{\sigma} c^2} A_{1\parallel}^2 + \frac{\mu}{B} A_{1\parallel} \nabla_{\perp}^2 A_{1\parallel} \right) \end{aligned} \quad (79)$$

$$+ \frac{1}{2} \alpha \epsilon_{\delta}^2 \int d\Omega f_{\text{eq},e} \frac{e^2}{m_e c^2} A_{1\parallel}^2 + \alpha \epsilon_{\delta}^2 \int dV \frac{dV}{8\pi} |\nabla_{\perp} A_{1\parallel}|^2. \quad (80)$$

For the Padé approximated model, the expression for the field energy is

$$\begin{aligned} \mathcal{E}_{\text{F}}^{\text{Padé}} = & \epsilon_{\delta} \frac{1}{2} q_i \int d\Omega (1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}) \langle \phi_1 \rangle f_i \\ & - \epsilon_{\delta} \frac{1}{2} e \int d\Omega (1 - \nabla_{\perp} \cdot \rho_i^2 \nabla_{\perp}) \phi_1 f_e. \end{aligned}$$

In the case of the model with adiabatic and hybrid electrons, the expressions for the conserved energy have to be discussed separately since they are derived from a slightly different variational formulation, which combines a fluid and kinetic formalism. With adiabatic electrons, the corresponding contribution to the energy should be considered as a field term:

$$\begin{aligned} \mathcal{E} = & \sum_{\sigma \neq e} \int d\Omega (H_0 + \epsilon_{\delta} H_1) f_{\sigma} + \epsilon_{\delta}^2 \sum_{\sigma \neq e} \int d\Omega H_2 f_{\text{eq},\sigma} \\ & + \epsilon_{\delta} \int dV \left[n_{e0} \phi_1 + \epsilon_{\delta} \frac{e}{2T_e} n_{e0} (\phi_1 - \bar{\phi}_1)^2 \right], \end{aligned} \quad (81)$$

where the last term is considered as a field term that includes the energy of the adiabatic electrons in the system. Following the

general procedure, we substitute the test function $\hat{\phi}_1 = \phi_1$ into the quasineutrality equation. The field energy is then given by

$$\mathcal{E}_F = \epsilon_\delta \frac{1}{2} \sum_{\sigma \neq e} \int d\Omega q_\sigma f_\sigma \langle \phi_1 \rangle + \epsilon_\delta \frac{1}{2} \int dV n_{e0} \phi_1 \quad (82)$$

and the kinetic part of energy consists of the ion contribution only:

$$\mathcal{E}_{\text{kin}} = \sum_{\sigma \neq e} \int d\Omega f_\sigma H_0 = \sum_{\sigma \neq e} \int d\Omega f_\sigma \left(\frac{m_\sigma v_\parallel^2}{2} + \mu B \right). \quad (83)$$

3. Numerical implementation

ORB5 uses a low-noise δf PIC method [64,65] consisting of separating the full distribution function f into a prescribed, time-independent background distribution f_0 and a perturbed, time-dependent distribution δf such that only the latter is discretized using markers, or numerical particles, that are used to sample the phase space. Furthermore, the code uses an operator splitting approach which consists of solving first for the collisionless dynamics and then considering the collisions and various sources. The time integration of the collisionless dynamics is made using a 4th-order Runge–Kutta (RK4). The collisions are treated with a Langevin approach.

This section describes the numerical implementation of the gyrokinetic equations presented in the previous section. First, the low-noise δf PIC method as well as the field discretization and solving are presented. Then, the noise reduction techniques, essential to control the noise inherent to the finite sampling of phase space with particles, are described. Finally, the different heat sources, relevant diagnostics, and the parallelization of the code are discussed. In this section, we omit the subscripts s specifying the species for the sake of simplifying the notation.

3.1. Discretization of δf and equations of motion

In ORB5, phase space is sampled using a set of N markers that are distributed according to a function $g(z, t)$ which is discretized as

$$g(z, t) \simeq \sum_{i=1}^N \frac{\delta[z - z_i(t)]}{J(z)}, \quad (84)$$

where $\delta[x]$ is the Dirac distribution, z is a set of generalized phase-space coordinates, $z_i(t)$ is the orbit of the i th marker in phase space, and $J(z)$ is the Jacobian associated with the coordinates z . Even though the choice of the distribution function $g(z, t)$ is not constrained, we make the convenient choice of using a distribution satisfying

$$\frac{dg}{dt}(z, t) = 0, \quad (85)$$

where the d/dt operator is the collisionless total time derivative defined by the general Vlasov equation, Eq. (14). In other words, Eq. (85) is ensured by evolving the trajectories $z_i(t)$ of the markers along the same characteristics as the physical particles. Both background and perturbed distribution functions can be linked to the marker distribution by the weight fields $W(z, t)$ and $P(z, t)$:

$$\begin{aligned} f_0 &= P(z, t)g(z, t) \simeq P(z, t) \sum_{i=1}^N \frac{\delta[z - z_i(t)]}{J(z)} \\ &= \sum_{i=1}^N P(z_i(t), t) \frac{\delta[z - z_i(t)]}{J(z)} = \sum_{i=1}^N p_i(t) \frac{\delta[z - z_i(t)]}{J(z)}, \end{aligned} \quad (86)$$

$$\begin{aligned} \delta f &= W(z, t)g(z, t) \simeq W(z, t) \sum_{i=1}^N \frac{\delta[z - z_i(t)]}{J(z)} \\ &= \sum_{i=1}^N W(z_i(t), t) \frac{\delta[z - z_i(t)]}{J(z)} = \sum_{i=1}^N \delta w_i(t) \frac{\delta[z - z_i(t)]}{J(z)}, \end{aligned} \quad (87)$$

where $p_i(t) = P(z_i(t), t)$ and $\delta w_i(t) = W(z_i(t), t)$ are the marker weights representing respectively the amplitude of f_0 and δf carried by each marker. The total distribution functions are normalized such that

$$\int f(z, t)J(z)dz = N_{\text{ph}}, \quad (88)$$

where N_{ph} is the physical number of particles in the system. Note that the coefficient N_{ph}/N is hereafter included in the weights such that $p_i(t) \equiv (N_{\text{ph}}/N)p_i(t)$ and $\delta w_i(t) \equiv (N_{\text{ph}}/N)\delta w_i(t)$.

3.1.1. Solving for the collisionless dynamics

According to the time splitting approach, the collisionless dynamics is solved first using the standard δf or the direct δf [66] methods. For the standard δf the time evolution of a marker weight δw_i is given by

$$\begin{aligned} \frac{d}{dt}\delta w_i(t) &= \frac{d}{dt}W(z_i(t), t) = \frac{d}{dt} \left[\frac{\delta f(z, t)}{g(z, t)} \right] = \frac{1}{g(z, t)} \frac{d}{dt}\delta f(z, t) \\ &\quad - \frac{\delta f(z, t)}{g(z, t)^2} \frac{d}{dt}g(z, t). \end{aligned} \quad (89)$$

The last term cancels due to the choice for the evolution of the distribution function $g(z, t)$, Eq. (85). The total distribution function f being constant along collisionless trajectories in phase space, the evolution equation of δw_i , Eq. (89), can be written as

$$\frac{d}{dt}\delta w_i(t) = -\frac{1}{g(z, t)} \frac{d}{dt}f_0(z_i(t)) = -p_i(t) \frac{1}{f_0(z_i(t))} \frac{d}{dt}f_0(z_i(t)). \quad (90)$$

Similarly, an equation for the p_i weight can also be derived:

$$\frac{d}{dt}p_i(t) = \frac{1}{g(z, t)} \frac{d}{dt}f_0(z_i(t)) = p_i(t) \frac{1}{f_0(z_i(t))} \frac{d}{dt}f_0(z_i(t)). \quad (91)$$

In ORB5, both weight equations as well as the numerical particle trajectories are integrated in time using an RK4 method.

On the other hand, the direct δf method exploits the invariance of the total distribution function f along the nonlinear collisionless trajectories; this property is not ensured in the linear and/or neoclassical limits. It allows one to directly evaluate the weights without numerically solving a differential equation. Adding Eqs. (90) and (91) leads to

$$\begin{aligned} \frac{d}{dt}(\delta w_i(t) + p_i(t)) &= 0 \implies \delta w_i(t) + p_i(t) \\ &= \delta w_i(t_0) + p_i(t_0), \quad \forall t, \end{aligned} \quad (92)$$

which comes from the invariance of both the f and g distribution functions. Furthermore, rewriting Eq. (91), we find

$$\frac{d}{dt} \left[\ln \left(\frac{p_i(t)}{f_0(z_i(t))} \right) \right] = 0 \implies \frac{p_i(t)}{f_0(z_i(t))} = \frac{p_i(t_0)}{f_0(z_i(t_0))}. \quad (93)$$

The direct δf algorithm thus consists of first evaluating the $p_i(t)$ weight according to Eq. (93) and then computing the $\delta w_i(t)$ weight using Eq. (92). Note that whatever the δf method used, if the collisionless limit is considered only the δw_i weights are required since the distribution $g(z, t)$ is invariant along the marker trajectories. Indeed, inserting Eq. (93) into Eq. (90) gives

$$\frac{d}{dt}\delta w_i(t) = -\frac{p_i(t_0)}{f_0(z_i(t_0))} \frac{d}{dt}f_0(z_i(t)), \quad (94)$$

which, in the limit $\Delta t \rightarrow 0$, gives

$$\delta w_i(t + \Delta t) \simeq \delta w_i(t) - \frac{1}{g(z_i(t_0))} [f_0(z_i(t + \Delta t)) - f_0(z_i(t))]. \quad (95)$$

Therefore, in the collisionless limit, we do not need to explicitly evolve $p_i(t)$.

3.1.2. Solving for the collisional dynamics

The collision operators are derived assuming linearization with respect to a local Maxwellian distribution. However, ORB5 is typically operated using the canonical background Maxwellian distribution as control variates in order to keep the background distribution in equilibrium in the collisionless gyrokinetic equation. Upon entering the collisions module, the δw_i and p_i weights are therefore converted to represent the perturbation with respect to a local Maxwellian background distribution, and are reverted when leaving it [27]. In this section, f_0 and δf always refer to these converted distributions, i.e. $f_0 = f_L$ and $\delta f = f - f_L$. At each time step, the collision operators are applied sequentially after the collisionless dynamics.

The electron-ion collision operator and the test-particle component of the intra-species collision operators are applied using a Langevin approach. In the gyrokinetic framework, this corresponds to applying random “kicks” in velocity space.

For electrons colliding on ions, Eq. (66) is reformulated in a spherical coordinate system in velocity space with radius v , polar angle θ , and azimuthal angle ζ in which the incoming electron's velocity corresponds to $\theta = 0$. Then, coming back to the ORB5 set of coordinates, the outgoing velocity of the electron is given by

$$v_{\parallel, \text{out}} = v_{\text{in}} \left[-\sin(\Delta\theta) \sin(\zeta_{\text{out}}) \sqrt{1 - \xi_{\text{in}}^2} + \xi_{\text{in}} \cos(\Delta\theta) \right], \quad (96)$$

$$v_{\perp, \text{out}}^2 = v_{\text{in}}^2 - v_{\parallel, \text{out}}^2, \quad (97)$$

where the $\parallel$ and $\perp$ subscripts are the components of the velocity parallel and perpendicular to the magnetic field, $\Delta\theta = 2R\sqrt{v_{\text{ei}}(v)}\Delta t$, R being a random sample of a PDF with mean 0 and variance 1, and ζ_{out} is a random sample of a uniform distribution between 0 and 2π . Note that the kinetic energy is exactly conserved by this procedure as in the original model.

Applying a similar approach for the “test-particle” self-collisions, Eqs. (59)–(62) yield the following outgoing particle trajectory:

$$v_{\parallel, \text{out}} = \frac{1}{v_{\text{in}}} \left[-\Delta v_y v_{\perp, \text{in}} + (v_{\text{in}} + \Delta v_z) v_{\parallel, \text{in}} \right], \quad (98)$$

$$v_{\perp, \text{out}}^2 = \Delta v_x^2 + \frac{1}{v_{\text{in}}^2} \left[\Delta v_y v_{\parallel, \text{in}} + (v_{\text{in}} + \Delta v_z) v_{\perp, \text{in}} \right]^2, \quad (99)$$

where Δv_x , Δv_y , and Δv_z are the particle change in velocity, with the unit vector $\hat{z}$ in the direction of the incoming particle's velocity. These kicks are described by

$$\Delta v_x = v_{\text{th}} \sqrt{\frac{K(v_{\text{in}}) \bar{v} \Delta t}{2}} R_1, \quad (100)$$

$$\Delta v_y = v_{\text{th}} \sqrt{\frac{K(v_{\text{in}}) \bar{v} \Delta t}{2}} R_2, \quad (101)$$

$$\Delta v_z = -H(v_{\text{in}}) v_{\text{in}} \bar{v} \Delta t + v_{\text{th}} \sqrt{H(v_{\text{in}}) \bar{v} \Delta t} R_3, \quad (102)$$

where R_1 , R_2 , and R_3 are again independent random numbers sampled from a PDF with mean 0 and variance 1. The marker magnetic moment is then updated accordingly.

It can be shown [27] that the evolution of the marker weight δw_i due to collisions can be expressed as

$$\frac{d}{dt} \delta w_i(t) = -p_i(t) \frac{C[\delta f, f_L]}{f_L} \Big|_{[z_i(t), t]}, \quad (103)$$

where $z_i(t)$ is the marker position after the “test-particle” kicks. At this point, the “background-reaction” operator, Eq. (63), is slightly corrected as follows:

$$\Delta \delta w_i(t) = -\frac{p_i}{n_b} \left[(1 - 3\sqrt{\pi} G(x)) \Delta N_b + 6\sqrt{\pi} H(x) \frac{\Delta P_{\parallel, b} v_{\parallel, \text{out}, r}}{v_{\text{th}, b}^2} + \sqrt{\pi} G(x) \frac{\Delta E_b}{v_{\text{th}, b}^2} \right], \quad (104)$$

where ΔN_b , $\Delta P_{\parallel, b}$ and ΔE_b are determined by imposing that the combined effect of applying the test particle collision operator and the background reaction term conserves mass, parallel momentum and kinetic energy within each configuration space bin b to machine precision.

3.1.3. Particle loading

At the beginning of a simulation, the markers are loaded in phase space using a Halton–Hammersley sequence [67,68] and according to the distribution function $g(z, t = 0) = f_{\text{rad}}(s) f_v(v_{\parallel}, v_{\perp})$, where $f_{\text{rad}}(s)$ and $f_v(v_{\parallel}, v_{\perp})$ define respectively the radial and velocity sampling distributions. In ORB5, the spatial sampling is defined by the *specified loading* distribution function $f_{\text{rad}}(s) = 1 - f_g + f_g \exp[(s - s_0)^2 / \Delta s^2]$, where $f_g \in [0, 1]$, s_0 , and Δs are input parameters. In velocity space $(v_{\parallel}, v_{\perp})$, the distribution is either uniform in the half-plane $(v_{\parallel}, v_{\perp} > 0)$ within $v = \sqrt{v_{\parallel}^2 + v_{\perp}^2} \leq \kappa_v v_{\text{th}, \sigma}$ with

$$f_v(v_{\parallel}, v_{\perp}) = \frac{1}{v_{\perp} (\pi \kappa_v v_{\text{th}}(s))^2}, \quad (105)$$

or uniform in 3D velocity space within a velocity sphere of radius $\kappa_v v_{\text{th}, \sigma}$ with

$$f_v(v_{\parallel}, v_{\perp}) = \frac{1}{\frac{4}{3} \pi \kappa_v^3 v_{\text{th}}^3(s)}, \quad (106)$$

where κ_v is an input parameter usually set at $\kappa_v = 5$ and which allows one to define a cut-off in velocity sampling.

For the marker weight initialization, two main schemes are implemented. The first option is a *white noise initialization* defined by

$$\delta w_i(t_0) = A(2Q_i - 1)p_i(t_0), \quad (107)$$

where Q_i is a quasi-random number in $[0, 1]$ given by the i th term of a van der Corput sequence [69] and A the maximum amplitude given as an input parameter, typically of the order of $A \sim 10^{-3} - 10^{-5}$. The disadvantage of this scheme is that the initial density perturbation is inversely proportional to the number of particles and the time until physical modes emerge from the initial state is roughly proportional to the number of particles. To accelerate the mode development, the *mode initialization* can be used. It consists in initializing a number of Fourier modes:

$$\delta w_i(t_0) = \frac{A_0 p_i(t_0)}{(m_2 - m_1 + 1)(n_2 - n_1 + 1)} \left| \frac{T(s_0)}{\nabla T(s_0)} \right| \times \left| \frac{T(s_i(t_0))}{\nabla T(s_i(t_0))} \right| \sum_{m=m_1}^{m_2} \sum_{n=n_1}^{n_2} \cos(m\theta_i^*(t_0) - n\varphi_i(t_0)), \quad (108)$$

where A_0 , n_1 , n_2 , m_1 , m_2 are input parameters. Typically, for linear simulations of microinstabilities with a toroidal mode number n_0 , it is convenient to use $n_1 = n_2 = n_0$ and $m_1 = m_2 = -n_0 q(s_0)$ as modes are almost aligned with the magnetic field lines. Finally, whatever initialization is used, the initial average value of the weights is set to zero:

$$\frac{1}{N} \sum_i^N \delta w_i(t_0) = 0. \quad (109)$$

As mentioned in Section 2.1 the markers are pushed in toroidal magnetic coordinates (s, θ^*, φ) . To avoid the singularity that would appear in the equations of motion at the magnetic axis, the coordinate system is changed to $(\xi, \eta, \varphi) = (s \cos \theta^*, s \sin \theta^*, \varphi)$ near the axis. All equilibrium quantities for both *ad-hoc* and MHD equilibria are loaded on an (R, Z) , where R and Z are cylindrical coordinates, grid and are linearly interpolated to an (s, θ^*) grid. Markers that exit the radial domain at $s > 1$ are reflected back into the plasma at a position which conserves toroidal momentum, the particle energy, and the magnetic moment but with a null weight to avoid unphysical accumulation of perturbed density at the radial edge.

3.2. Quasineutrality and Ampère equations

In ORB5 the quasineutrality and Ampère equations are solved using the Galerkin method considering linear, quadratic, or cubic B-splines finite elements defined on a $(N_s, N_{\theta^*}, N_\varphi)$ grid. The perturbed fields ϕ and $A_\parallel$ hereafter noted $\Psi = \{\phi, A_\parallel\}$ are thus discretized as follows:

$$\Psi(\mathbf{X}, t) = \sum_{\mu} \Psi_{\mu}(t) \Lambda_{\mu}(\mathbf{X}), \quad (110)$$

where $\{\Psi_{\mu}(t)\}$ are the field coefficients and $\{\Lambda_{\mu}(\mathbf{X})\}$ are a tensor product of 1D B-splines of degree $p = \{1, 2, 3\}$, $\Lambda_{\mu}(\mathbf{X}) = \Lambda_j^p(s) \Lambda_k^p(\theta^*) \Lambda_l^p(\varphi)$, with $\mu = (j, k, l)$.

Using the decomposition defined in Eqs. (87) and (110), and setting the test functions $\hat{\phi}_1 = \Lambda_v(\mathbf{X})$, $v = (j', k', l')$ of the variational forms of the quasineutrality and Ampère equations, Eqs. (9) and (13), leads to a linear system of the form

$$\sum_{\mu} A_{\mu\nu} \Psi_{\mu}(t) = b_{\nu}(t), \quad (111)$$

where $A_{\mu\nu}$ and b_{ν} are respectively a real symmetric positive-definite square matrix and a column vector that are defined by the physical models used in the quasineutrality and Ampère equations. Due to the finite support of the B-splines, the matrix $A_{\mu\nu}$ is usually a block diagonal matrix composed of banded submatrices. Note that in the case of the quasi-neutrality valid to all orders, the matrix A in general becomes a full matrix. As illustration, we show here the linear system for the case of a single ion species plasma in the limit of adiabatic electrons with the long wavelength approximation for the ion polarization density:

$$A_{\mu\nu}^{\text{LWA, adiab}} = \int \left[\frac{en_0(\psi)}{T_e(\psi)} (\Lambda_{\mu}(\mathbf{X}) \Lambda_{\nu}(\mathbf{X}) - \bar{\Lambda}_{\mu}(s) \bar{\Lambda}_{\nu}(s)) + \frac{n_0(\psi) m_i}{B^2} \nabla_{\perp} \Lambda_{\mu}(\mathbf{X}) \cdot \nabla_{\perp} \Lambda_{\nu}(\mathbf{X}) \right] dV, \quad (112)$$

$$b_{\nu}(t) = \sum_{p=1}^N \frac{\delta w_p(t)}{2\pi} \int_0^{2\pi} d\Theta \Lambda_{\nu}(\mathbf{X}_p + \rho_{L,p}(\Theta)), \quad (113)$$

where $\rho_{L,p}$ is the Larmor radius of a particle p . Here, the gradient perpendicular to the magnetic field is approximated by the gradient in the poloidal plane, i.e. $\nabla_{\perp} \simeq \nabla_{\text{pol}} = \nabla s \frac{\partial}{\partial s} + \nabla \theta^* \frac{\partial}{\partial \theta^*}$. Note that the expression for $b_{\nu}(t)$, Eq. (113), is independent of the choice of coordinates. This is due to the particle representation of δf , Eq. (87), and the Galerkin finite element method based on the variational form of the field equations, Eqs. (9) and (13). This is very convenient practically as the charge deposition is totally transparent to the choice of the coordinates system, which greatly simplifies the numerical implementation. A more complete description of the discretized Poisson equation for arbitrary wavelengths can be found in [30].

In ORB5, the linear system of equations, Eq. (111), is represented in Fourier space to obtain an equivalent system of equations [70] that is solved using the FFTW library [71] and a direct solver from the LAPACK library [72]. This Fourier representation of the fields is convenient because it allows one to decouple the toroidal mode numbers n , provided that the field equations are linear and that the unperturbed system is axisymmetric. Furthermore, the modes of interest, e.g. drift-wave type and Alfvén waves, are typically almost aligned with the magnetic field lines and can be described with just a small set of Fourier coefficients, which greatly decreases the numerical cost as compared to solving the system in direct space. Noting $\mathcal{F}$ the double discrete Fourier transform on both poloidal and toroidal directions, the linear system of Eqs. (111) becomes

$$\sum_{\mu} \mathcal{F} A_{\mu\nu} \mathcal{F}^{-1} \mathcal{F} \Psi_{\mu} = \mathcal{F} b_{\nu}, \quad (114)$$

$$\mathcal{F} A_{\mu\nu} \mathcal{F}^{-1} = \hat{A}_{(j,j')^{(n,m),(n',m')}}, \quad (115)$$

$$\mathcal{F} \Psi_{\mu} = \hat{\Psi}_j^{n,m}, \quad (116)$$

$$\mathcal{F} b_{\nu} = \hat{b}_{j'}^{n,m'}, \quad (117)$$

where n and m are respectively the toroidal and poloidal Fourier mode numbers.

Due to the axisymmetry of the system and because the Poisson–Ampère system of equation is linear, the toroidal direction can be decoupled from the others with $n = n'$ [25]:

$$\sum_j \sum_m \hat{A}_{(j,j')^{(n,m),(n',m')}} \hat{\Psi}_j^{n,m} = \frac{\hat{b}_{j'}^{n,m'}}{M^{n,p}} \quad \forall n, \quad (118)$$

where the matrix $M^{n,p}$ is defined by

$$M^{n,p} = \sum_{l'=1}^{N_{\varphi}} \int d\varphi \Lambda_{l'}^p(\varphi) \Lambda_{l'}^p(\varphi) \exp \left[\frac{2\pi i}{N_{\varphi}} n(l' - l) \right], \quad (119)$$

and can be computed analytically for any B-spline of order p :

$$M^{n,1} = \frac{2\pi}{N_{\varphi}} \left[\frac{2}{3} + \frac{1}{3} \cos \left(\frac{2\pi}{N_{\varphi}} n \right) \right], \quad (120)$$

$$M^{n,2} = \frac{2\pi}{N_{\varphi}} \left[\frac{8}{15} + \frac{13}{30} \cos \left(\frac{2\pi}{N_{\varphi}} n \right) + \frac{1}{30} \cos^2 \left(\frac{2\pi}{N_{\varphi}} n \right) \right], \quad (121)$$

$$M^{n,3} = \frac{2\pi}{N_{\varphi}} \left[\frac{136}{315} + \frac{33}{70} \cos \left(\frac{2\pi}{N_{\varphi}} n \right) + \frac{2}{21} \cos^2 \left(\frac{2\pi}{N_{\varphi}} n \right) + \frac{1}{630} \cos^3 \left(\frac{2\pi}{N_{\varphi}} n \right) \right]. \quad (122)$$

This toroidal decoupling is very interesting numerically because the different toroidal mode numbers n can be solved for separately.

The matrix $A_{\mu\nu}$ and the right-hand side b_{ν} are modified such that the following boundary conditions are used. At the magnetic axis the *unicity condition* is applied, $\Psi(s = 0, \theta^*, \varphi, t) = \Psi(s = 0, \theta^* = 0, \varphi, t)$, $\forall \theta^*$. At the outer radial edge, Dirichlet boundary conditions are applied, $\Psi(s = 1, \theta^*, \varphi, t) = 0$. Note that ORB5 can also be run in an annulus, i.e. $s \in [s_{\min}, s_{\max}]$, with $s_{\min} > 0$ and $s_{\max} < 1$, in which case Dirichlet boundary conditions are applied on both edges. For the quasineutrality equation with polarization density valid to all orders in $k_{\perp} \rho$, the equation is integral and no Dirichlet boundary conditions need to be applied [30].

3.2.1. Gyroaveraging

For all gyroaveraging operations, the plane of the Larmor ring is approximated to lie in the poloidal plane. The number of gyro-points can be either fixed or determined by an adaptive scheme;

in which case a fixed number of Larmor points is used for all the particles having a Larmor radius smaller or equal to the thermal Larmor radius and the number of points increases linearly for larger Larmor radii. Usually, a fixed number of 4 gyropoints is sufficient for perturbations up to $k_{\perp}\rho_L \sim 1$. However, using the adaptive scheme reduces the noise as it acts as a Bessel filter averaging out shorter wavelength fluctuations [73].

In magnetic coordinates, the positions of the gyropoints are parametrized using the gyroangle Θ :

$$\mathbf{x}(\Theta) = \mathbf{X} + \boldsymbol{\rho}(\Theta) = \mathbf{X} + \rho \frac{\nabla S}{|\nabla S|} \cos \Theta + \rho \frac{\mathbf{b} \times \nabla S}{|\mathbf{b} \times \nabla S|} \sin \Theta, \quad (123)$$

where $\mathbf{X}$ is the position of the guiding center.

The gradients of gyroaveraged electric potential, $\nabla \langle \phi_1 \rangle$, are approximated as

$$\nabla_{\mathbf{X}} \langle \phi_1 \rangle \simeq \langle \nabla \phi_1 \rangle = \frac{1}{2\pi} \oint_0^{2\pi} \nabla_{\mathbf{X}} \phi_1(\mathbf{X} + \boldsymbol{\rho}) d\Theta, \quad (124)$$

where the subscript $\mathbf{X}$ stands for the gradient with respect to the gyrocenter coordinates and Θ is the gyroangle. We define a new set of coordinates $\bar{\mathbf{X}} = (\bar{R}, \bar{Z}) = (R + \rho \cos \Theta, Z + \rho \sin \Theta) = \mathbf{X} + \boldsymbol{\rho}$ representing the position of the particle on the gyro-ring in the poloidal plane where $\bar{R}$ is in the direction of the major axis and $\bar{Z}$ is in the direction of the vertical axis. This coordinate system is the projection of Eq. (123) on the (R, Z) plane of a cylindrical coordinate system. Using the chain rule, the term $\nabla_{\mathbf{X}} \phi_1$ from Eq. (123) can be written as

$$\nabla_{\mathbf{X}} \phi_1(\mathbf{X} + \boldsymbol{\rho}) = \nabla_{\bar{\mathbf{X}}} \phi_1 - \frac{\rho}{2} \left(\frac{\partial \phi_1}{\partial \bar{R}} \cos \Theta + \frac{\partial \phi_1}{\partial \bar{Z}} \sin \Theta \right) \frac{\nabla_{\mathbf{X}} B}{B}. \quad (125)$$

A similar procedure is done for $\nabla \langle A_{1\parallel} \rangle$. In ORB5, Eq. (125) can be either directly evaluated as in [36] or approximated by neglecting the second term, leading to $\nabla_{\mathbf{X}} \phi_1 \approx \nabla_{\bar{\mathbf{X}}} \phi_1$.

3.2.2. Fourier filter

Typical physical modes of interest, e.g. drift waves and low-frequency Alfvén waves, are mainly aligned with the magnetic field lines, i.e. they have $m \approx nq(s)$. Due to this strong anisotropy, only a limited set of (n, m) Fourier coefficients is required to describe the physical modes as the amplitude of the (n, m) Fourier coefficients rapidly decreases away from $m = nq(s)$ [9]. It is then beneficial to filter out all the non physically relevant Fourier modes in order to reduce the sampling noise and maximize the timestep size. The filter is applied on the Fourier coefficients of the perturbed density and current to remove all the non physical Fourier modes introduced by the charge and current depositions:

$$\tilde{b}_{(j,k,l)} = \sum_{n,m} f_{j,n,m} \hat{b}_j^{n,m} e^{im\theta_k^*} e^{in\varphi_l}, \quad (126)$$

where $f_{j,n,m}$ is the Fourier filter that in general depends on the radius (radial index j), and the poloidal and toroidal mode numbers m and n , respectively.

Two different filters are used successively. First, a *rectangular filter*, which is the simplest one, is applied such that all the modes outside of the window $[n_{\min}, n_{\max}] \times [m_{\min}, m_{\max}]$ specified in input are filtered out. This filter is not sufficient as it keeps modes with $k_{\parallel}/k_{\perp}$ much bigger than ρ^* , which is inconsistent with the gyrokinetic ordering [9]. Since the modes of interest are mainly aligned with the magnetic field, i.e. they satisfy $k_{\parallel}\rho_i \simeq [m - nq(s)][q(s)R]^{-1}\rho_i = \mathcal{O}(\rho^*)$, a second surface-dependent *field-aligned filter* is applied. It consists in retaining only m modes close to $nq(s)$, i.e. $m \in [nq(s) - \Delta m, nq(s) + \Delta m]$, where Δm is an

input parameter specifying the width of the filter. With this field-aligned filter, the maximum value of $|k_{\parallel}|$ represented is $|k_{\parallel}|_{\max} = |\Delta m|/qR$. Since $|k_{\parallel}|_{\max}\rho_i$ scales with ρ^* , the value of Δm required to describe all physically relevant modes is invariant with the system size. Typically, a value of $\Delta m = 5$ is sufficient [70]. In summary, for each mode $n \in [n_{\min}, n_{\max}]$ only the modes $m \in [m_{\min}, m_{\max}] \cap [-nq(s) \pm \Delta m]$ are retained.

3.3. Noise control techniques

Due to the finite number of markers used to sample phase space, PIC simulations are subject to sampling errors, referred to as numerical noise, which in fact increases in a nonlinear simulation, thus deteriorating the signal quality and forbidding long simulations without noise control techniques. All the difficulty of such noise-reducing schemes is to actually control the weight growth without creating severe non-physical artifacts. In this section we present the different noise control schemes implemented in ORB5.

3.3.1. Krook operator

The Krook operator implemented in ORB5 [31] is a source term which weakly damps the non axisymmetric fluctuations without significantly affecting the zonal flows. This is done via a correction term that also allows one to conserve various moments by projecting out some components of the source. The Krook noise-control term, S_K^{NC} , is composed of a relaxation term and its correction S_K^{corr} :

$$S_K^{\text{NC}} = -\gamma_K \delta f + S_K^{\text{corr}}, \quad (127)$$

$$S_K^{\text{corr}} = \sum_{i=1}^{N_{\text{mom}}} g_i(s) M_i f_0, \quad (128)$$

where γ_K is the Krook damping rate. The correction term is a sum over the N_{mom} moments M_i one wishes to conserve on a flux-surface average. Typically, in ORB5, the moments that can be conserved are the density, parallel velocity, zonal flows, and kinetic energy. They are respectively defined by $M_i = \{1, v_{\parallel}, v_{\parallel}/B - (v_{\parallel}/B), \mathcal{E}_K\}$, where the tilde represents the bounce average and $\mathcal{E}_K$ is the kinetic energy of a particle. Note that, as shown in [31], the conservation of the density and $v_{\parallel}/B - (v_{\parallel}/B)$ moment is there to ensure conservation of the zonal flow long wavelength residual computed in [74]. The coefficients $g_i(s)$ are defined such that there is no contribution of the source to a given moment M_j , i.e.

$$\int dW \bar{M}_j S_K^{\text{NC}} = 0, \quad \forall j, \quad (129)$$

where the bar notation stands for the flux-surface average. Injecting the definition of the Krook source term, Eq. (127), in Eq. (129) leads to a linear system of equations that is solved at each time step and radial position to find the coefficients $g_i(s)$:

$$\sum_i^{N_{\text{mom}}} S_{ij}(s) g_i(s) = \delta S_j(s), \quad (130)$$

with

$$S_{ij}(s) = \int dW \bar{M}_j(\mathbf{X}, v_{\parallel}, \mu) M_i(\mathbf{X}, v_{\parallel}, \mu) f_0(\mathbf{X}, v_{\parallel}, \mu), \quad (131)$$

$$\delta S_j(s) = \gamma_K \int dW \delta f(\mathbf{X}, v_{\parallel}, \mu, t) M_j(\mathbf{X}, v_{\parallel}, \mu). \quad (132)$$

Note that the flux-surface average is numerically represented by a binning of the markers in the radial direction. This implies that the conservation is ensured only on average across each radial bin.

As already mentioned, the noise control should not affect significantly the turbulence. To this end, values of the order of one tenth of the maximum linear growth rate are usually used for the Krook damping rate γ_K . In this way, the linear phase is not substantially modified and a high signal-to-noise ratio can be obtained. On the other hand, applying this noise control technique is not appropriate when considering collisions as the damping rate γ_K is comparable to typical values of the collision frequency thus interfering with the effect of collisions.

By construction, the Krook operator damps the fluctuations to restore the full distribution function to its initial state. If the kinetic energy is not conserved while conserving the other moments, it allows one to run temperature gradient-driven simulations by acting as an auto-regulated heat source while allowing for the unconstrained evolution of the density and flow profiles.

3.3.2. Coarse-graining

Coarse-graining [63,75] is an additional noise-control method implemented in ORB5 [38] to reduce the problems of weight-spreading and filamentation of the distribution function, that lead to large mean squared particle weights. The idea is essentially to dissipate fine-scale structures of the distribution function in phase space, as represented by the marker weights. This is an improvement compared to the Krook operator, which only preserves certain moments of the zonal distribution function but otherwise somewhat indiscriminately damps the whole distribution function; the Krook operator, for example, was found to be unsuitable for neoclassical studies for which collisional effects are clearly essential.

In a Eulerian code, phase space dissipation is often implemented as a hyper-viscosity operator on the grid in the five spatial and velocity directions. As the PIC approach does not involve a phase-space grid, we need an alternative method to smooth the weights of nearby markers.

Computationally, the method consists of binning the particles in phase space cells which are field-aligned in configuration space, and then reducing the deviation of particle weights in the grid cell from their average value. To avoid smoothing structures at the turbulence scale too strongly, the bins must be small compared to typical length and velocity scales; on the other hand the bins need to frequently contain more than one marker for this procedure to be effective. Elongated field-aligned bins are used because the distribution function varies much more rapidly perpendicular to the field line than along it.

The bins are volumes in a block-structured Cartesian mesh in coordinates $(s, z, \theta^*, \lambda, \epsilon)$, with the number of bins uniform in each direction, except that the number of bins in the θ^* direction is proportional to s , so that the spatial volume of bins is roughly constant. The coordinate ϵ is the particle kinetic energy, λ is the pitch angle, and z is a field-line label that is computed as

$$z = \varphi - q(s)[\theta^* - \theta_0^*(\theta^*)], \quad (133)$$

with θ_0^* the center of the bin in the θ^* direction:

$$\theta_0^*(\theta^*) = \left(\left\lfloor \frac{\theta^* + \pi}{\Delta\theta^*} \right\rfloor + \frac{1}{2} \right) \Delta\theta^* - \pi, \quad (134)$$

where $\lfloor \bullet \rfloor$ is the floor function and $\Delta\theta^*$ is the width of a bin in the θ^* direction. With this choice of z we have a field-aligned bin, but we also have $z \sim \varphi$ if there are many bins in the θ^* direction because $\theta_0^*(\theta^*)$ tends to θ^* for an infinite number of bins. This is useful because the domain decomposition – discussed in detail in Section 3.5 – means that markers on a single processor have a small range of values of φ . Thus, the first step in the binning process is to distribute the markers according to z and move them to this alternative domain decomposition. In the z decomposition,

coarse-graining is local to each domain, so we do not need to communicate quantities on the 5D coarse-graining mesh.

The number of bins in the s and z directions is the field mesh quantities N_s and N_φ respectively and the number of θ^* , energy and pitch-angle bins is specified as input parameters. To avoid excessive damping of zonal flows, around 32 bins are needed in each of the energy and pitch-angle directions. Often 16 bins in the θ^* direction are sufficient to avoid excessive damping of parallel structures.

The smoothing operation changes the particle weight w by an amount $\mathcal{N}\Delta t\gamma_{cg}(\bar{w} - w)$, where $\bar{w}$ is the average particle weight in the bin, $\mathcal{N}$ is the number of timesteps (of length Δt) between coarse-graining operations, and γ_{cg} is a parameter controlling the coarse-graining rate. In the limit of large number of markers, this leads to a damping of fine-scale structures in the distribution function with a rate γ_{cg} . Note, however, that in practice, typical runs have 0.1 markers per bin, so that the effective coarse-graining rate is lower than γ_{cg} by a factor of 10.

3.3.3. Quad-tree particle-weight smoothing

The grid-based coarse-graining procedure has the possible drawback of being inaccurate if the grid of phase-space bins is too fine so that the local statistics is not good enough, or being very diffusive if the grid is too coarse. An alternative procedure, gridless in velocity space and more probabilistic in nature, has been proposed in [39] and implemented in ORB5. It consists in pairing neighboring markers and replacing their weights by an average, weighted by a function of their distance in velocity space. The way of computing the distance and the weight has an influence on the diffusivity of the method, for this reason we use a procedure for pairing only particles which are close enough. Since the gyrokinetic velocity space is 2D, the pairing procedure is done using a quad-tree algorithm: first, the particles are binned in the configuration space and then, a quad-tree procedure is applied to define regions in velocity space within which particles will be paired. This works by subdividing recursively the 2D velocity space in four sub-boxes until the number of particles in a sub-box is smaller than a given value set as an input parameter. At this point, the particles within a sub-box are randomly paired and their weight is changed according to the following procedure: for a pair of two markers with weights δw_1 and δw_2 and velocities $\mathbf{v}_1$ and $\mathbf{v}_2$, the new weights are given by

$$w_1^{\text{new}} = (1 - \Gamma) w_1^{\text{old}} + \Gamma \bar{w}, \quad (135)$$

$$w_2^{\text{new}} = (1 - \Gamma) w_2^{\text{old}} + \Gamma \bar{w}, \quad (136)$$

with

$$\Gamma = e^{-\frac{(\mathbf{v}_1^x - \mathbf{v}_2^x)^2 + (\mathbf{v}_1^y - \mathbf{v}_2^y)^2}{h_v^2}}, \quad (137)$$

$$\bar{w} = \frac{w_1^{\text{old}} + w_2^{\text{old}}}{2}, \quad (138)$$

where the x and y superscripts are used to identify the two dimensions of the velocity and the h_v parameter defines how strong is the smoothing procedure with respect to the distance separating the pair of markers in velocity space. Note that, by construction, the smoothing operation conserves the total weight, ensuring density conservation. By picking different pairs of particles within the same quad-tree sub-box, the smoothing operation can be applied several times per timestep. Typically, one smoothing step is done at every timestep.

3.3.4. Enhanced control variates

The ORB5 code solves the uncoupled electromagnetic gyrokinetic equations in the p_z -formulation, Eqs. (9) and (13) and therefore includes the cancellation problem [76] which, if untreated, in

practice limits the electromagnetic simulations to very-low-beta cases, $\beta < \sqrt{m_e/m_i}$, where β is the stored kinetic energy divided by the magnetic field energy. Different methods mitigating this problem have been developed for the particle-in-cell framework in Refs. [35,36,77–79] and for the Eulerian approach in Ref. [18]. In ORB5 the cancellation problem is treated [34] using the enhanced control variates scheme presented in [35,36]. A further development of the mitigation schemes is given in Refs. [80,81], the so-called pullback mitigation based on the mixed-variable formulation [82] of the gyrokinetic theory, has also been implemented in ORB5 [37]. Mitigation of the cancellation problem has made possible the ORB5 electromagnetic simulations described in Refs. [83,84].

The enhanced control variates approach is based on the decomposition of the distribution function into the so-called adiabatic and nonadiabatic parts introduced in [85] while constructing a perturbative procedure for the solution to the gyrokinetic Vlasov equation. The same decomposition can be extracted via the pull-back transformation between the particle distribution function and the reduced gyrokinetic distribution [6]. This transformation requires that the equilibrium distribution commutes with the background dynamics, i.e. $\{f_{eq}, H_0\} = 0$. Furthermore, in ORB5 the distribution function is assumed to be a canonical Maxwellian, i.e. satisfying

$$\frac{\partial f_{eq,\sigma}}{\partial H_0} = -\frac{f_{eq,\sigma}}{T_\sigma}, \quad (139)$$

where the temperature is defined as

$$T_\sigma = \frac{1}{n_0} \int dW \frac{p_z^2}{2m_\sigma} f_{eq,\sigma}. \quad (140)$$

In the enhanced-control-variates scheme, the perturbed distribution function is split according to

$$f_{\sigma,1} = G_\sigma - \frac{f_{eq,\sigma}}{T_\sigma} H_1, \quad (141)$$

where the first and second terms are respectively the nonadiabatic and adiabatic parts.

The cancellation problem is related to the coexistence of very large and very small quantities in the variational form of the Ampère equation (13). To illustrate the problem, let us consider a case with only one ion species and rewrite Eq. (13). Note that the second and third integrals of Eq. (13) are the projections $\mathcal{J}_{s,\parallel}$ of the ion and electron currents onto the basis function $\hat{A}_{1\parallel}$:

$$\langle \mathcal{J}_{i,\parallel} \rangle \equiv \int dV \langle j_{i,\parallel} \rangle \hat{A}_{1\parallel} = \int d\Omega f_\sigma \frac{q_\sigma p_z}{m_\sigma c} \langle \hat{A}_{1\parallel} \rangle, \quad (142)$$

$$\mathcal{J}_{e,\parallel} \equiv \int dV j_{e,\parallel} \hat{A}_{1\parallel} = \int d\Omega f_e \frac{e p_z}{m_e c} \hat{A}_{1\parallel}. \quad (143)$$

Then, the fourth integral and the first term of the fifth integral of Eq. (13) are the so-called skin terms and can be written as

$$\int d\Omega f_{eq,\sigma} \left(\frac{4\pi q_\sigma^2}{m_\sigma c^2} A_{1\parallel} \hat{A}_{1\parallel} \right) = \frac{\beta_\sigma}{\rho_{th,s}^2} \int dV A_{1\parallel} \hat{A}_{1\parallel}, \quad (144)$$

where one defines $\beta_\sigma = 4\pi n_\sigma T_\sigma / B^2$ and $\rho_{th,s}$ is the thermal Larmor radius. Finally, the remaining terms of Eq. (13) are combined to form

$$\begin{aligned} & \int dV \nabla_\perp A_{1\parallel} \cdot \nabla_\perp \hat{A}_{1\parallel} + \int d\Omega f_{eq,i} \frac{2\pi\mu}{B} \\ & \times [A_{1\parallel} \nabla_\perp^2 \hat{A}_{1\parallel} + \hat{A}_{1\parallel} \nabla_\perp^2 A_{1\parallel}] \\ & = \int dV \nabla_\perp \cdot [(1 - \beta_i) \nabla_\perp A_{1\parallel} \hat{A}_{1\parallel}], \end{aligned} \quad (145)$$

where the integration by parts has been used and the terms containing second order gradients of the background quantities neglected. Putting Eqs. (142)–(145) back into Eq. (13) leads to

$$\begin{aligned} & \frac{\beta_i}{\rho_{th,i}^2} \int d\Omega f_{eq,i} A_{1\parallel} \hat{A}_{1\parallel} + \frac{\beta_e}{\rho_{th,e}^2} \int d\Omega f_{eq,e} A_{1\parallel} \hat{A}_{1\parallel} \\ & - \int d\Omega \nabla_\perp \cdot [(1 - \beta_i) \nabla_\perp A_{1\parallel} \hat{A}_{1\parallel}] = \frac{4\pi}{c} (\langle \mathcal{J}_{i,\parallel} \rangle - \mathcal{J}_{e,\parallel}). \end{aligned} \quad (146)$$

The two skin terms can become very large, especially for electrons, due to their small mass. They cancel up to the second order FLR corrections with the adiabatic part of the currents $\langle j_{i,\parallel} \rangle$ and $j_{e,\parallel}$. This can be seen by splitting the currents into an adiabatic and nonadiabatic part using the splitting defined in Eq. (141) and injecting them back into Eq. (146).

The cancellation problem occurs in PIC simulations due to the different discretizations of particles and fields: the currents are typically computed using the particles while the skin terms are computed using the finite element grid. The terms to be canceled are much larger in magnitude than the remaining terms which are supposed to represent the physics. Therefore, the cancellation must be numerically extremely accurate, otherwise the relevant signal is dominated by numerical noise.

In ORB5, the cancellation problem is mitigated by discretizing both the skin terms and the adiabatic part of the currents in Eq. (146) with the same markers. The polarization-current term, $-\int d\Omega \nabla_\perp \cdot [(1 - \beta_i) \nabla_\perp A_{1\parallel} \hat{A}_{1\parallel}]$, is discretized on the grid since it does not contribute to the cancellation. This approach to the discretization is used in ORB5 in Ampère's law.

Ampère's law, Eq. (146), is used to compute the parallel magnetic potential $A_{1\parallel}$. Note that the non-adiabatic perturbed distribution function G_σ depends on $A_{1\parallel}$ which is unknown at this point of the computation. The solution is to use an easy-to-compute estimator, $\hat{S}$, and solve iteratively for $A_{1\parallel}$. In ORB5, the skin term $(\beta_s/\rho_s^2)A_{1\parallel}$ is used as a simple estimator for the $A_{1\parallel}$ -dependent part of the distribution function. One reformulates Ampère's law using the estimator $\hat{S}$:

$$(\hat{S} + L)a = (j - Sa) + \hat{S}a, \quad (147)$$

where a is the discretized magnetic vector potential component, S and L are respectively the discretized skin terms and polarization current term, and j represents the sum over the species of the discretized currents. For a good estimator, a small parameter $\|\hat{S} - S\| = \mathcal{O}(\epsilon)$ can be introduced to expand the vector potential, $a = a_0 + \epsilon a_1 + \epsilon^2 a_2 + \mathcal{O}(\epsilon^3)$. Ampère's law is then solved iteratively order by order in ϵ :

$$(\hat{S} + L)a_0 = j,$$

$$(\hat{S} + L)a_1 = (\hat{S} - S)a_0,$$

...

In practice, for typical production runs, less than 10 iterations are necessary. In ORB5, the estimator is expressed using the finite elements $\hat{S}_{kl} = \int \beta_\sigma / \rho_s^2 \Lambda_k(\mathbf{x}) \Lambda_l(\mathbf{x}) d^3x$. The marker-dependent part of the right-hand side of the iterative scheme is written as the enhanced control variates:

$$j_k - S_{kl} a_l^{n-1} = \sum_{v=1}^{N_p} p_{zv} \left(\delta w_v + \frac{q_\sigma p_z}{m_\sigma c} \frac{\langle A_{1\parallel}^{(n-1)} \rangle}{T_\sigma} f_{eq,\sigma}(Z_v) \zeta_v \right) \langle \Lambda_k \rangle_v, \quad (148)$$

where ζ_v is the elementary phase-space volume associated with a given marker. The same enhanced control variates are used also for the perturbed particle density. In practice, it results in a straightforward and computationally cheap modification of




Fig. 1. MPI parallelization using domain decomposition in the toroidal direction and domain cloning.

the charge and current assignment routines in ORB5. Since the computations are mainly dominated by particle operations, the field solve only takes a few percent of the total timing of ORB5. With this iterative scheme turned on, the total timing increases by a few percent only.

3.4. Heating operators

A primary goal of simulating the full plasma core (by contrast to a local approach) is to examine the self-consistent evolution of plasma profiles in the presence of both turbulence-driven transport and external sources. In practice, even for running global simulations where realistic global profile evolution is not of interest, it is generally inconvenient to run simulations without a heat source: if the goal is to look at transport properties at a specific temperature gradient, simulations where the temperature gradient relaxes rapidly evolve away from the desired parameters. In ORB5 temperature gradient control and injection of energy flux are imposed through sources added to the r.h.s. of the Vlasov equation. These do not model the detailed physics of a realistic heat source (for example, the radially localized temperature anisotropy generated by resonant heating schemes) but simply control moments of the distribution function.

For the control of the temperature gradient, so that it stays close to an initial gradient, a *thermal relaxation operator* is used (this can be seen as an effective interaction with a heat bath) of the form

$$S_{H1} = -\gamma_H \left[\delta f(\epsilon, s) - f_0(\epsilon, s) \frac{\int d^3v \delta f(\epsilon, s)}{\int d^3v f_0(\epsilon, s)} \right]. \quad (149)$$

Contrarily to Eq. (127), here δf depends only on the radial position s and the kinetic energy ϵ . This source term maintains the distribution function $f(\epsilon, s)$ close to the initial value, i.e. it relaxes back to f_0 with a rate γ_H . Note that the heating operator, Eq. (149), does not act as a noise control, unlike the modified Krook operator defined in Section 3.3.1. Indeed, this operator applies the same correction to all the markers within a velocity bin, which does not reduce the noise caused by filamentation in velocity space. The second term in the equation ensures that the gyrocenter density is not modified by the source term, i.e. the heat source does not act as an effective particle source. Due to the symmetry of this operator in $v_{||}$ it also does not add parallel momentum to the system; testing [31,86] has shown that long wavelength flows are largely unaffected by this heat source although certain higher order effects could lead to significant flow drive on shorter wavelengths [42].

The choice of γ_H determines how strongly the temperature gradient is clamped to the initial gradient; since the form of the heating is not physical, it is necessary to set γ_H small enough to not excessively damp temperature corrugations; empirical investigations suggest that setting γ_H ten times smaller than typical instability growth rates is appropriate to ensure convergence. It is possible to specify this heat source to be active only in certain regions of the plasma, so that, for example, a “source-free” region in the middle of the simulation domain may be obtained.

Fixed-input power simulations may be obtained by using a *fixed heat source* of the form

$$S_{H2} = \gamma_R(s) \frac{\partial f_0}{\partial T}, \quad (150)$$

where $\gamma_R(s)$ defines the spatial heating profile written in terms of an effective inverse timescale over which the local temperature would vary in the absence of transport. Generally this operator is used to represent a fixed input power source in the core of the tokamak. To model the energy losses near the edge, two options can be chosen: first, to define a fixed heat sink by setting a profile $\gamma_R(s)$ with negative values in the edge region; second, to define a buffer region near the boundary in which a Krook operator is specified (see Section 3.3.1), thus damping the edge profiles close to their initial values.

3.5. Parallelization

In order to simulate complex physical systems in a reasonable amount of time, the ORB5 code is massively parallelized using a hybrid MPI/OpenMP and MPI/OpenACC implementation. The MPI parallelization is done using both domain cloning and domain decomposition [87,88] techniques, Fig. 1.

The physical domain is first replicated into disjoint clones and the markers are evenly distributed among them. Each clone can be further decomposed by splitting the physical domain in the toroidal direction into subdomains. Each subdomain of each clone is attributed to an MPI task such that the total number of processes is given by $P_{MPI} = P_{sub} \times P_{clones}$, where P_{sub} and P_{clones} are respectively the number of tasks attributed to the subdomains and clones.

After each time step, data must be transferred between the clones and subdomains. For the clones, mainly global reductions of grid quantities are required, e.g. after each charge deposition step all the contributions from the clones must be gathered to compute the self-consistent electromagnetic fields. For the subdomains, it consists of nearest neighbor communications for the guard cells, global communications of grid data (parallel data




Fig. 2. Strong scalings on the Piz Daint machine (Cray XC50, 12 cores per node). Speed-up is defined as inverse ratio of wall clock time to single node wall clock time, multiplied by powers of 8 for the weak scaling factor. The number of particles indicates the number of ions and the number of electrons.

transpose) for Fourier transforms and point to point communications of particle data where we exchange the particles that have moved from a subdomain to another. Note that in ORB5, the particle exchange algorithm is not restricted to the nearest neighbors, all-to-all is supported.

While the domain decomposition scales well with the number of subdomains, a large number of clones is problematic in terms of performance. Indeed, the domain cloning approach is quickly limited by the more demanding communications and the memory congestion due to the field data replication. To overcome this issue each MPI task is multithreaded using OpenMP. This has the main advantage of limiting the number of clones while still increasing the code performance by sharing the workload among threads.

To take advantage of the new HPC platforms equipped with accelerators, the ORB5 code has been recently ported to GPU using OpenACC. These developments will be detailed in a separate paper [89]. The choice of using OpenMP and OpenACC was motivated because they allow us to keep all parallelization options in a single source code version.

4. Results

4.1. Parallel scalability

In Fig. 2, we perform series of strong scalings of a typical electromagnetic simulation with kinetic electrons. All the runs are made on the Piz Daint supercomputer hosted at CSCS in Switzerland which is a hybrid Cray XC40/XC50 machine. For this scaling, up to 4096 compute nodes of the XC50 partition equipped with one 12-core Intel Xeon E5-2690 v3 at 2.60 GHz are used.

We use as many ions as electrons, using an adaptive number of Larmor points per guiding center going from 4 to 32. The simulations are nonlinear, with a fixed number of 2 iterations for the control variate scheme. Cubic splines are used. Scalar and 1D diagnostics are computed every other time step and 2D diagnostics one step out of ten.

The starting point of each strong scaling makes a weak scaling where the grid resolution is multiplied by 2 in each dimension, the number of particles by 8 and the number of compute nodes by 8. We use domain cloning inside nodes and domain decomposition in between them, meaning that the number of clones is set to the number of cores per node, i.e. 12, and the number of subdomains to the number of nodes. We make an exception for the large-scale cases where the number of nodes exceeds the number of toroidal cells, i.e. the last points of the $640 \cdot 10^6$

particles and $5120 \cdot 10^6$ particles cases, in which case we double the number of clones so that the number of parallel tasks is equal to the product of subdomains and number of clones.

ORB5 scales very well up to 128 nodes with a speed-up larger than 85% of the ideal speed-up. We even get a small superscaling from 2 to 16 nodes thanks to increased data locality and decreased memory congestion.

Above 256 nodes, the speed-up is limited mainly by the MPI communications of parallel data transpose required for the field Fourier transforms. Some effort is currently put on reducing the cost of those communications.

The performance of the GPU-accelerated ORB5 will be assessed in a following paper. In short, for scaling tests similar to Fig. 2, representative of production runs, the GPU-accelerated ORB5 is up to 4 times faster than the CPU-only code.

4.2. Strong flows and toroidal rotation

We demonstrate the use of the strong flow features of the code using an adiabatic electron CYCLONE benchmark case with nominal toroidal rotation rate $\Omega_R = 0.2c_s/R$. The numerical parameters are similar to those used for typical global CYCLONE benchmark cases with sources [41] (circular concentric equilibrium, $\rho^* = 1/180$, $a/R = 0.36$, plateau-like initial logarithmic temperature gradient profiles with $R/L_T = 6.9$ and $R/L_n = 2.2$). The field solver grid is $N_s \times N_{\theta^*} \times N_{\varphi} = 128 \times 512 \times 256$, and 1.2×10^8 markers are used. A heating operator is used with the rate $0.013c_s/a$ to maintain temperature profiles near their initial value. Coarse graining is applied every $2.8a/c_s$ time units, with 64 bins in energy and pitch angle, and a blending factor of 1 (so all weights in a coarse-graining bin are set equal).

The effects of strong rotation on the equilibrium have been discussed earlier for ORB5 [33], so we focus on demonstrating the operation of the code in the nonlinear regime; at the moderate levels of rotation tested here the effects are not expected to be dramatic. As in non-rotating simulations, there is some overall relaxation of the heat profiles as the turbulence driven transport commences, Figs. 3–4. The parallel flow profile, Fig. 5, is not constrained by the heating operator and relaxes slightly (note that the initial parallel velocity profile is not completely flat, as might be expected for solid body rotation). In these simulations, although strong flow effects due to centrifugal and Coriolis drift are included, the pinch driven momentum flux is expected to be nearly zero due to the use of an adiabatic electron model: this is consistent with the observation of little net momentum flux in these simulations [90].




Fig. 3. Temperature gradient R/L_T versus time and radius in a strong flow simulation.




Fig. 4. Ion heat flux in gyro-Bohm units versus time and radius in a strong flow simulation.




Fig. 5. Zonal average of parallel velocity versus radius at the beginning (red dashed trace) and end (blue trace) of a strong flow simulation.

4.3. GK simulations of Alfvén modes in the presence of turbulence

One of the main recent developments of ORB5 has been to allow for electromagnetic simulations. The electromagnetic extension, via the control-variate scheme [35,80] was initially implemented in 2011 [26] and proved to work for ITG instabilities. Further improvements have been necessary for having successful shear-Alfvén wave (SAW) tests, and the first results have been published in 2016 [83]. SAWs are known to be crucial in present tokamaks and future reactors, as they can be driven unstable by energetic particles (EP), which can deteriorate the EP confinement [5].

The linear dynamics of Alfvén instabilities investigated with ORB5 has been recently benchmarked against analytical theory and other codes [47,83]. Moreover, the nonlinear dynamics of Alfvén modes due to the wave-particle nonlinearity has been investigated with ORB5, and compared with the GK code EUTERPE [84,91], where in particular a detailed study of the saturation levels due to wave-particle nonlinearity has been carried out. Finally, after a dedicated phase of verification and benchmarking on the Alfvén dynamics, ORB5 has now started the investigation of the self-consistent interaction of Alfvén instabilities and turbulence. Here, we describe a test case where the nonlinear dynamics of an Alfvén mode is investigated in the presence of turbulence.

The tokamak geometry and magnetic field is taken consistently with Ref. [83], for the case referred to as energetic particle modes. Regarding the bulk profiles, the ion and electron temperatures are taken equal everywhere, $T_e(s) = T_i(s)$. Here, differently from Ref. [83], a value of $T_e(s = \text{speak})$ corresponding to $\rho^* = \rho_s/a = 0.00571$, is chosen. The electron thermal to magnetic pressure ratio is $\beta_e = 5 \cdot 10^{-4}$. An analytical function is used for the profiles of the equilibrium density and temperature, for the three species of interest (thermal deuterium, labeled here as “d”, thermal electrons, labeled here as “e”, and hot deuterium, labeled here as “EP”). For the EP density, for example, the function is written as $n_{EP}(s)/n_{EP}(s_r) = \exp\{-\Delta \kappa_n \tanh[(s - s_r)/\Delta]\}$. The value of Δ is the same for all species, for both density and temperature: $\Delta = 0.208$. Deuterium and electrons have $\kappa_n = 0.3$ and $\kappa_T = 1.0$, and the EP have $\kappa_n = 10.0$ and $\kappa_T = 0.0$. The EP temperature is given by $T_{EP}/T_e = 100$. The distribution function of the EP population is Maxwellian in $p_{||}$. The EP averaged concentration is $\langle n_{EP} \rangle / n_e = 0.002$. A filter allows poloidal and toroidal mode numbers with $-128 < m < 128$ and $0 \leq n < 40$ to develop. Unicity boundary conditions are imposed at $s = 0.0$ and Dirichlet at $s = 1.0$. A white noise initial perturbation is set at $t = 0$. The electron mass is chosen as $m_e/m_i = 0.005$. A Krook operator is applied to deuterium and electrons.

Nonlinear collisionless electromagnetic simulations have been performed with ORB5, with turbulence driven by the equilibrium temperature gradients, peaked at mid-radius with and without EP. In the absence of EP, heat transport exhibits radial corrugated structures, Fig. 6; larger corrugations are observed in the inner half of the radial domain, i.e. $s \in [0, 0.5]$. Those corrugations are also present as long-lived structures in the $E \times B$ velocity profile, Fig. 7 (left), and are particularly visible for $s \in [0, 0.5]$. Avalanches of $E \times B$ velocity are generated at $s \sim 0.5$ with a frequency matching the local GAM frequency, Fig. 7 (right), and then propagate outward with constant frequency. Finally, the nonzonal component of the scalar potential has been measured, and observed to grow linearly in the first so-called linear phase of the ITG turbulence, and then saturate.

In the simulation where EPs are loaded, an Alfvénic instability is observed growing on top of the turbulence [93,94]. Zonal structures, like zero frequency zonal flows and geodesic acoustic modes, play the role of mediators of small-scale turbulence and large-scale Alfvén modes. A comparison of the perturbed electric potential with and without EP is shown in Fig. 8.




Fig. 6. Corrugations of the effective heat diffusivity radial profile (top). The heat diffusivity is in gyro-Bohm units, $\chi_{GB} = \rho_e^2 c_s / a$. Radial profile of the temperature logarithmic gradient R/L_T (bottom). All the profiles are time averaged in $t \in [500, 1515] [a/c_s]$.

Such simulations are numerically demanding due to the fact that they investigate intrinsically multi-scale phenomena. Thus, high resolution in space and time is needed like in turbulence simulations, the electrons must be treated kinetically for driving the current perturbations necessary for the evolution of the Alfvén physics, and three separate plasma species (thermal ions, thermal electrons, and EP) must be evolved simultaneously, in order to drive the Alfvén mode unstable, with a corresponding high number of markers.

5. Conclusion

ORB5 is a global PIC code used to solve the electromagnetic gyrokinetic equations in presence of collisions and various sources, e.g. heat and strong flows. The Vlasov–Maxwell model on which it relies is derived from variational principles and all the different

physical approximations are included in the gyrokinetic action. This allows to consistently derive the equations of motion while ensuring conservation properties that can be later used to assess the simulation quality for example.

Three models are available for the Poisson equation in which the ion polarization density can be represented at full order, up to the second order in FLR corrections, or using a Padé approximation. On the other hand, Ampère's equation is computed up to second order in FLR corrections. While the ions are a gyrokinetic species, the electrons can be treated as adiabatic, drift-kinetic, or a hybrid mix where passing electrons are adiabatic and trapped electrons are drift-kinetic. Furthermore, the hybrid model, which does not respect the ambipolarity condition, has been corrected.

The code is based on the PIC δf control variate scheme in order to reduce the numerical noise due to finite particle sampling and various other techniques are used to further limit this noise, e.g. noise reduction schemes are implemented to constrain the particle weight spreading and a Fourier filter allows to solve only the physically relevant modes. The ORB5 code is parallelized using a hybrid OpenMP/MPI or OpenACC/MPI approach allowing to benefit from the many and multicore HPC systems. Scalability experiments have shown ORB5's excellent parallel scalability up to thousands of cores. A few physical simulations run with the ORB5 code and including e.g. strong flows, toroidal rotation, and shear Alfvén waves are also presented to illustrate the capabilities of the code.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors deeply acknowledge past valuable contributions of Dr. T. M. Tran to the ORB5 code development. Fruitful discussions with S. Ethier are gratefully acknowledged.

The authors thank the CSCS for providing access to the full Piz Daint machine to perform the scalability tests. This work was supported by a grant from the Swiss National Supercomputing Centre (CSCS) under project ID s760. This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014–2018 and 2019–2020 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission. This work was partly supported by the Swiss National Science Foundation.




Fig. 7. $E \times B$ velocity as a function of time and radius (left) and the corresponding frequency spectrum (right). The white dashed line represents analytical estimates of the GAM frequency [92].




Fig. 8. Poloidal cut of the perturbed potential $\phi - \bar{\phi}$. On the left, a poloidal cut showing the ITG turbulence without energetic particles. On the right, the characteristic poloidal structure of the reversed-shear Alfvén eigenmode ($n = 1, m = 2$) in the presence of turbulence is shown.

References

- [1] R.J. Fonck, G. Cosby, R.D. Durst, S.F. Paul, N. Bretz, S. Scott, E. Synakowski, G. Taylor, *Phys. Rev. Lett.* 70 (24) (1993) 3736–3739, <http://dx.doi.org/10.1103/PhysRevLett.70.3736>, URL <https://link.aps.org/doi/10.1103/PhysRevLett.70.3736>.
- [2] S.J. Zweben, J.A. Boedo, O. Grulke, C. Hidalgo, B. LaBombard, R.J. Maqueda, P. Scarin, J.L. Terry, *Plasma Phys. Control. Fusion* 49 (7) (2007) S1–S23, <http://dx.doi.org/10.1088/0741-3335/49/7/S01>, URL <http://stacks.iop.org/0741-3335/49/i=7/a=S01?key=crossref.b88a2f06af2f78dc5d83138198588e28>.
- [3] W. Horton, *Rev. Modern Phys.* 71 (3) (1999) 735–778, <http://dx.doi.org/10.1103/RevModPhys.71.735>, URL <http://link.aps.org/doi/10.1103/RevModPhys.71.735>.
- [4] C. Cheng, L. Chen, M. Chance, *Ann. Phys. (N. Y.)* 161 (1) (1985) 21–47, [http://dx.doi.org/10.1016/0003-4916\(85\)90335-5](http://dx.doi.org/10.1016/0003-4916(85)90335-5), URL <https://www.sciencedirect.com/science/article/pii/0003491685903355?via%3Dihub>.
- [5] L. Chen, F. Zonca, *Rev. Modern Phys.* 88 (1) (2016) 015008, <http://dx.doi.org/10.1103/RevModPhys.88.015008>, URL <https://link.aps.org/doi/10.1103/RevModPhys.88.015008>.
- [6] A.J. Brizard, T. Hahm, *Rev. Modern Phys.* 79 (2) (2007) 421–468, <http://dx.doi.org/10.1103/RevModPhys.79.421>, URL <http://link.aps.org/doi/10.1103/RevModPhys.79.421>, <http://journals.aps.org/rmp/abstract/10.1103/RevModPhys.79.421>.
- [7] N. Tronko, C. Chandre, J. Plasma Phys. 84 (3) (2017) <http://dx.doi.org/10.1017/s0022377818000430>, arXiv:1709.05222. URL <http://www.cambridge.org/core/journals/journal-of-plasma-physics/article/second-order-nonlinear-gyrokinetic-theory-from-the-particle-to-the-gyrocenter/C377BD7B6F0E11D8EB6E3D92AD97C7FE> <http://arxiv.org/abs/1709.05222>.
- [8] X. Garbet, Y. Idomura, L. Villard, T.H. Watanabe, *Nucl. Fusion* 50 (4) (2010) 043002, <http://dx.doi.org/10.1088/0029-5515/50/4/043002>, URL <http://stacks.iop.org/0029-5515/50/i=4/a=043002?key=crossref.c0e169b0cfa8ae95a37226a468dc019b>.
- [9] S. Jolliet, A. Bottino, P. Angelino, R. Hatzky, T. Tran, B.F.B. McMillan, O. Sauter, K. Appert, Y. Idomura, L. Villard, *Comput. Phys. Comm.* 177 (5) (2007) 409–425, <http://dx.doi.org/10.1016/j.cpc.2007.04.006>, URL <http://linkinghub.elsevier.com/retrieve/pii/S0010465507002251>, <http://www.sciencedirect.com/science/article/pii/S0010465507002251>.
- [10] Y. Idomura, S. Tokuda, Y. Kishimoto, *Nucl. Fusion* 43 (4) (2003) 234–243, <http://dx.doi.org/10.1088/0029-5515/43/4/303>, URL <http://iopscience.iop.org/article/10.1088/0029-5515/43/4/303>, <http://stacks.iop.org/0029-5515/43/i=4/a=303?key=crossref.d169703164fcc8d437ddd6f6a74be33>.
- [11] Z. Lin, T.S. Hahm, W.W. Lee, W.M. Tang, R.B. White, *Science* (80-) 281 (5384) (1998) 1835–1837, <http://dx.doi.org/10.1126/science.281.5384.1835>, URL <http://www.ncbi.nlm.nih.gov/pubmed/9743492>.
- [12] S. Ku, C.S. Chang, R. Hager, R.M. Churchill, G.R. Tynan, I. Cziegler, M. Greenwald, J. Hughes, S.E. Parker, M.F. Adams, E. D’Azevedo, P. Worley, *Phys. Plasmas* 25 (5) (2018) 056107, <http://dx.doi.org/10.1063/1.5020792>, URL <http://aip.scitation.org/doi/10.1063/1.5020792>.
- [13] W.X. Wang, T.S. Hahm, W.W. Lee, G. Rewoldt, J. Manickam, W.M. Tang, *Phys. Plasmas* 14 (7) (2007) 072306, <http://dx.doi.org/10.1063/1.2750647>, URL <http://aip.scitation.org/doi/10.1063/1.2750647>.
- [14] A.M. Dimits, T.J. Williams, J.A. Byers, B.I. Cohen, *Phys. Rev. Lett.* 77 (1) (1996) 71–74, <http://dx.doi.org/10.1103/PhysRevLett.77.71>, URL <https://link.aps.org/doi/10.1103/PhysRevLett.77.71>.
- [15] T. Korpilo, A. Gurichenko, E. Gusakov, J. Heikkinen, S. Janhunen, T. Kiviniemi, S. Leerink, P. Niskala, A. Perevalov, *Comput. Phys. Comm.* 203 (2016) 128–137, <http://dx.doi.org/10.1016/j.cpc.2016.02.021>, URL <https://www.sciencedirect.com/science/article/pii/S0010465516300364?via%3Dihub>.
- [16] F. Jenko, W. Dorland, M. Kotschenreuther, B.N. Rogers, *Phys. Plasmas* 7 (5) (2000) 1904, <http://dx.doi.org/10.1063/1.874014>, URL <https://aip.scitation.org/doi/10.1063/1.874014>.
- [17] Y. Idomura, M. Ida, T. Kano, N. Aiba, S. Tokuda, *Comput. Phys. Comm.* 179 (6) (2008) 391–403, <http://dx.doi.org/10.1016/j.cpc.2008.04.005>, URL <https://www.sciencedirect.com/science/article/pii/S0010465508001409?via%3Dihub>.
- [18] J. Candy, R.E. Waltz, J. Comput. Phys. 186 (2) (2003) 545–581, [http://dx.doi.org/10.1016/S0021-9991\(03\)00079-2](http://dx.doi.org/10.1016/S0021-9991(03)00079-2), URL <https://www.sciencedirect.com/science/article/pii/S0021999103000792>, <https://www.sciencedirect.com/science/article/pii/S0021999103000792?via%3Dihub>.
- [19] A.G. Peeters, D. Strintzi, *Phys. Plasmas* 11 (8) (2004) 3748–3751, <http://dx.doi.org/10.1063/1.1762876>, URL <http://aip.scitation.org/doi/10.1063/1.1762876>.
- [20] T.-H.H. Watanabe, H. Sugama, *Nucl. Fusion* 46 (1) (2006) 24–32, <http://dx.doi.org/10.1088/0029-5515/46/1/003>, URL <http://stacks.iop.org/0029-5515/46/i=1/a=003?key=crossref.3d7e6f40ce52b566eca20514f822347d>.
- [21] M. Kotschenreuther, G. Rewoldt, W.M. Tang, *Comput. Phys. Comm.* 88 (2–3) (1995) 128–140, [http://dx.doi.org/10.1016/0010-4655\(95\)00035-E](http://dx.doi.org/10.1016/0010-4655(95)00035-E), URL <https://www.sciencedirect.com/science/article/pii/001046559500035E?via%3Dihub>.
- [22] V. Grandgirard, J. Abiteboul, J. Bigot, T. Cartier-Michaud, N. Crouseilles, G. Dif-Pradalier, C. Ehrlacher, D. Esteve, X. Garbet, P. Ghendrih, G. Latu, M. Mehrenberger, C. Norscini, C. Passeron, F. Rozar, Y. Sarazin, E. Sonnendrücker, A. Strugarek, D. Zarzoso, *Comput. Phys. Comm.* 207 (2016) 35–68, <http://dx.doi.org/10.1016/j.cpc.2016.05.007>, URL <http://linkinghub.elsevier.com/retrieve/pii/S0010465516301230>.
- [23] C.K. Birdsall, A.B. Langdon, *Plasma Physics via Computer Simulations*, CRC press, 2004, p. 504.
- [24] W. Lee, J. Comput. Phys. 72 (1) (1987) 243–269, [http://dx.doi.org/10.1016/0021-9991\(87\)90080-5](http://dx.doi.org/10.1016/0021-9991(87)90080-5), URL <http://www.sciencedirect.com/science/article/pii/0021999187900805>, <https://aip.scitation.org/doi/10.1063/1.864140>.
- [25] T.M. Tran, K. Appert, M. Fivaz, G. Jost, J. Vaclavik, L. Villard, *Theory of Fusion Plasmas*, Società Italiana di Fisica, 1999, p. 45.
- [26] A. Bottino, T. Vernay, B. Scott, S. Brunner, R. Hatzky, S. Jolliet, B.F. McMillan, T.M. Tran, L. Villard, *Plasma Phys. Control. Fusion* 53 (12) (2011) 124027, <http://dx.doi.org/10.1088/0741-3335/53/12/124027>, URL <http://stacks.iop.org/0741-3335/53/i=12/a=124027?key=crossref.7fb008935b961f69d91dc474edce4036>.

- [27] T. Vernay, S. Brunner, L. Villard, B.F. McMillan, S. Jolliet, T.M. Tran, A. Bottino, J.P. Graves, *Phys. Plasmas* 17 (12) (2010) 122301, <http://dx.doi.org/10.1063/1.3519513>, URL http://pop.aip.org/resource/1/phpaen/v17/i12/p122301_s1 <http://aip.scitation.org/doi/10.1063/1.3519513>.
- [28] S. Jolliet, Gyrokinetic Particle-in-Cell Global Simulations of Ion-Temperature-Gradient and Collisionless-Trapped-Electron-Mode Turbulence in Tokamaks, (Ph.D. thesis), École Polytechnique Fédérale de Lausanne, 2009, <http://dx.doi.org/10.5075/epfl-thesis-4326>, URL <https://infoscience.epfl.ch/record/130837?ln=fr>.
- [29] T. Vernay, S. Brunner, L. Villard, B.F. McMillan, S. Jolliet, A. Bottino, T. Görler, F. Jenko, *Plasma Phys. Control. Fusion* 55 (7) (2013) 074016, <http://dx.doi.org/10.1088/0741-3335/55/7/074016>, URL <http://stacks.iop.org/0741-3335/55/i=7/a=074016?key=crossref.6e26708d569dcb2dcd6631cc3a584de7>.
- [30] J. Dominski, B.F. McMillan, S. Brunner, G. Merlo, T.M. Tran, L. Villard, *Phys. Plasmas* 24 (2) (2017) <http://dx.doi.org/10.1063/1.4976120>, URL <http://aip.scitation.org/toc/php/24/2>.
- [31] B.F. McMillan, S. Jolliet, T.M. Tran, L. Villard, A. Bottino, P. Angelino, *Phys. Plasmas* 15 (5) (2008) 1–10, <http://dx.doi.org/10.1063/1.2921792>, URL <http://aip.scitation.org/doi/10.1063/1.2921792>.
- [32] S. Jolliet, B.F. McMillan, T. Vernay, L. Villard, A. Bottino, P. Angelino, *Phys. Plasmas* 16 (5) (2009) 52307–52310, <http://dx.doi.org/10.1063/1.3140036>, URL <http://link.aip.org/link/?PHP/16/52307/1>.
- [33] J.D. Collier, B.F. McMillan, J.R. Robinson, *Phys. Plasmas* 23 (11) (2016) 112503, <http://dx.doi.org/10.1063/1.4965900>, URL <http://aip.scitation.org/doi/10.1063/1.4965900>.
- [34] A. Mishchenko, A. Bottino, R. Hatzky, E. Sonnendrücker, R. Kleiber, A. Könies, *Phys. Plasmas* 24 (8) (2017) 081206, <http://dx.doi.org/10.1063/1.4997540>, URL <http://aip.scitation.org/doi/10.1063/1.4997540>.
- [35] R. Hatzky, A. Könies, A. Mishchenko, J. Comput. Phys. 225 (1) (2007) 568–590, <http://dx.doi.org/10.1016/j.jcp.2006.12.019>, URL <https://www.sciencedirect.com/science/article/pii/S0021999106006085>.
- [36] R. Hatzky, R. Kleiber, A. Könies, A. Mishchenko, M. Borchardt, A. Bottino, E. Sonnendrücker, *J. Plasma Phys.* 85 (1) (2019) 905850112, <http://dx.doi.org/10.1017/S0022377819000096>, URL https://www.cambridge.org/core/product/identifier/S0022377819000096/type/journal_article.
- [37] A. Mishchenko, A. Bottino, A. Biancalani, R. Hatzky, T. Hayward-Schneider, N. Ohana, E. Lanti, S. Brunner, L. Villard, M. Borchardt, R. Kleiber, A. Könies, *Comput. Phys. Comm.* (2018) [arXiv:1811.05346](https://arxiv.org/abs/1811.05346). doi: [arXiv:1811.05346](https://arxiv.org/abs/1811.05346). URL https://www.sciencedirect.com/science/article/pii/S0010465518304181?dgcid=rss_sd_all <http://arxiv.org/abs/1811.05346>.
- [38] T. Vernay, S. Brunner, L. Villard, B.F. McMillan, S. Jolliet, T.M. Tran, A. Bottino, *Phys. Plasmas* 19 (4) (2012) 042301, <http://dx.doi.org/10.1063/1.3699189>, URL <http://scitation.aip.org/content/aip/journal/pop/19/4/10.1063/1.3699189>.
- [39] E. Sonnendrücker, A. Wachter, R. Hatzky, R. Kleiber, J. Comput. Phys. 295 (2015) 402–419, <http://dx.doi.org/10.1016/j.jcp.2015.04.004>, URL <https://www.sciencedirect.com/science/article/pii/S0021999115002442?via%3Dihub>.
- [40] V. Grandgirard, Y. Sarazin, P. Angelino, A. Bottino, N. Crouseilles, G. Darmet, G. Dif-Pradalier, X. Garbet, P. Ghendrih, S. Jolliet, G. Latu, E. Sonnendrücker, L. Villard, *Plasma Phys. Control. Fusion* 49 (12B) (2007) 173–182, <http://dx.doi.org/10.1088/0741-3335/49/12B/S16>, URL <http://stacks.iop.org/0741-3335/49/i=12B/a=S16?key=crossref.64dedbea6ea22f74127d3f5c432604dd> <http://iopscience.iop.org/0741-3335/49/12B/S16>.
- [41] X. Lapillonne, B.F. McMillan, T. Görler, S. Brunner, T. Dannert, F. Jenko, F. Merz, L. Villard, *Phys. Plasmas* 17 (11) (2010) 112321, <http://dx.doi.org/10.1063/1.3518118>, URL <http://aip.scitation.org/doi/10.1063/1.3518118>.
- [42] Y. Sarazin, V. Grandgirard, J. Abiteboul, S.J. Allfrey, X. Garbet, P. Ghendrih, G. Latu, A. Strugarek, G. Dif-Pradalier, P. Diamond, S. Ku, C. Chang, B. McMillan, T. Tran, L. Villard, S. Jolliet, A. Bottino, P. Angelino, *Nucl. Fusion* 51 (10) (2011) 103023, <http://dx.doi.org/10.1088/0029-5515/51/10/103023>, URL <http://stacks.iop.org/0029-5515/51/i=10/a=103023?key=crossref.5a1644f0c67bf3302c4798ff98f6af2>.
- [43] B.F. McMillan, X. Lapillonne, S. Brunner, L. Villard, S. Jolliet, A. Bottino, T. Görler, F. Jenko, *Phys. Rev. Lett.* 105 (15) (2010) 155001, <http://dx.doi.org/10.1103/PhysRevLett.105.155001>, URL <http://link.aps.org/doi/10.1103/PhysRevLett.105.155001>.
- [44] T. Görler, N. Tronko, W.A. Hornsby, A. Bottino, R. Kleiber, C. Nordsieck, V. Grandgirard, F. Jenko, E. Sonnendrücker, *Phys. Plasmas* 23 (7) (2016) 072503, <http://dx.doi.org/10.1063/1.4954915>, URL <http://aip.scitation.org/doi/10.1063/1.4954915>.
- [45] N. Tronko, A. Bottino, T. Görler, E. Sonnendrücker, D. Told, L. Villard, *Phys. Plasmas* 24 (5) (2017) 056115, <http://dx.doi.org/10.1063/1.4982689>, [arXiv:1703.07582](https://arxiv.org/abs/1703.07582). URL <http://aip.scitation.org/doi/10.1063/1.4982689>.
- [46] A. Biancalani, A. Bottino, C. Ehrlacher, V. Grandgirard, G. Merlo, I. Novikau, Z. Qiu, E. Sonnendrücker, X. Garbet, T. Görler, S. Leerink, F. Palermo, D. Zarzoso, *Phys. Plasmas* 24 (6) (2017) 062512, <http://dx.doi.org/10.1063/1.4985571>, URL <http://aip.scitation.org/doi/10.1063/1.4985571>.
- [47] A. Könies, S. Briguglio, N. Gorelenkov, T. Fehér, M. Isaev, P. Lauber, A. Mishchenko, D.A. Spong, Y. Todo, W.A. Cooper, R. Hatzky, R. Kleiber, M. Borchardt, G. Vlad, A. Biancalani, A. Bottino, *Nucl. Fusion* 58 (12) (2018) 126027, <http://dx.doi.org/10.1088/1741-4326/aae4e6>, URL <http://stacks.iop.org/0029-5515/58/i=12/a=126027?key=crossref.732f2cf421546adceace9e18a1fa64a>.
- [48] G. Merlo, J. Dominski, A. Bhattacharjee, C.S. Chang, F. Jenko, S. Ku, E. Lanti, S. Parker, *Phys. Plasmas* 25 (6) (2018) 062308, <http://dx.doi.org/10.1063/1.5036563>, URL <http://aip.scitation.org/doi/10.1063/1.5036563>.
- [49] A. Di Siena, A. Biancalani, T. Görler, H. Doerk, I. Novikau, P. Lauber, A. Bottino, E. Poli, T.A.U. Team, *Nucl. Fusion* 58 (10) (2018) 106014, <http://dx.doi.org/10.1088/1741-4326/aad51d>, URL <http://stacks.iop.org/0029-5515/58/i=10/a=106014?key=crossref.dc39673986e17b8827eef8596adb27ab>.
- [50] H. Lütjens, A. Bondeson, O. Sauter, *Comput. Phys. Comm.* 97 (3) (1996) 219–260, [http://dx.doi.org/10.1016/0010-4655\(96\)00046-X](http://dx.doi.org/10.1016/0010-4655(96)00046-X), URL <https://www.sciencedirect.com/science/article/pii/001046559600046X>.
- [51] H. Sugama, *Phys. Plasmas* 7 (2) (2000) 466–480, <http://dx.doi.org/10.1063/1.873832>, URL <http://aip.scitation.org/doi/abs/10.1063/1.873832> <http://aip.scitation.org/doi/10.1063/1.873832>.
- [52] N. Tronko, A. Bottino, E. Sonnendrücker, *Phys. Plasmas* 23 (8) (2016) 082505, <http://dx.doi.org/10.1063/1.4960039>, [arXiv:1604.03538](https://arxiv.org/abs/1604.03538). URL <http://aip.scitation.org/doi/10.1063/1.4960039>.
- [53] N. Tronko, A. Bottino, C. Chandre, E. Sonnendrücker, *Plasma Phys. Control. Fusion* 59 (6) (2017) 064008, <http://dx.doi.org/10.1088/1361-6587/aa68af>, [arXiv:1703.05095](https://arxiv.org/abs/1703.05095). URL <http://stacks.iop.org/0741-3335/59/i=6/a=064008?key=crossref.0311bba6987f428be09915498fab4f3>.
- [54] E. Lanti, J. Dominski, S. Brunner, B.F. McMillan, L. Villard, J. Phys. Conf. Ser. 775 (1) (2016) 012006, <http://dx.doi.org/10.1088/1742-6596/775/1/012006>, URL <http://stacks.iop.org/1742-6596/775/i=1/a=012006?key=crossref.5970237ead440732086f3dc78b4903bc>.
- [55] Y. Idomura, J. Comput. Phys. 313 (2016) 511–531, <http://dx.doi.org/10.1016/j.jcp.2016.02.057>.
- [56] A.Y. Aydemir, *Phys. Plasmas* 1 (4) (1994) 822, <http://dx.doi.org/10.1063/1.870740>, URL <http://scitation.aip.org/content/aip/journal/pop/1/4/10.1063/1.870740>.
- [57] P. Angelino, A. Bottino, R. Hatzky, S. Jolliet, O. Sauter, T.M. Tran, L. Villard, *Phys. Plasmas* 13 (5) (2006) 052304, <http://dx.doi.org/10.1063/1.2193947>, URL <http://aip.scitation.org/doi/10.1063/1.2193947> <http://scitation.aip.org/content/aip/journal/pop/13/5/10.1063/1.2193947>.
- [58] T.S. Hahm, *Phys. Plasmas* 3 (12) (1996) 4658–4664, <http://dx.doi.org/10.1063/1.872034>, URL <https://aip.scitation.org/doi/10.1063/1.872034> <http://aip.scitation.org/doi/10.1063/1.872034>.
- [59] L. Guazzotto, R. Betti, J. Manickam, S. Kaye, *Phys. Plasmas* 11 (2) (2004) 604–614, <http://dx.doi.org/10.1063/1.1637918>, URL <http://aip.scitation.org/doi/10.1063/1.1637918>.
- [60] B.F. McMillan, T. Vernay, A. Bottino, S. Jolliet, L. Villard, 37th EPS Conf. Plasma Phys. 2010, EPS 2010, Vol. 1, 2010, pp. 453–456.
- [61] G. Dif-Pradalier, V. Grandgirard, Y. Sarazin, X. Garbet, P. Ghendrih, *Phys. Rev. Lett.* 103 (6) (2009) 065002, <http://dx.doi.org/10.1103/PhysRevLett.103.065002>, URL <https://link.aps.org/doi/10.1103/PhysRevLett.103.065002>.
- [62] Z. Lin, W.M. Tang, W.W. Lee, *Phys. Plasmas* 2 (8) (1995) 2975–2988, <http://dx.doi.org/10.1063/1.871196>, URL <http://aip.scitation.org/doi/10.1063/1.871196>.
- [63] S. Brunner, E. Valeo, J.A. Krommes, *Phys. Plasmas* 6 (12) (1999) 4504–4521, <http://dx.doi.org/10.1063/1.873738>, URL http://pop.aip.org/resource/1/phpaen/v6/i12/p4504_s1%5Cnhttp://scitation.aip.org/getpdf/servlet/GetPDFServlet?filetype=pdf&id=PHPAEN000006000012004504000001&idtype=cvips&doi=10.1063/1.873738&prog=normal.
- [64] S.E. Parker, W.W. Lee, *Phys. Fluids B* 5 (1) (1993) 77–86, <http://dx.doi.org/10.1063/1.860870>, URL <http://aip.scitation.org/doi/10.1063/1.860870>.
- [65] G. Hu, J.A. Krommes, *Phys. Plasmas* 1 (4) (1994) 863–874, <http://dx.doi.org/10.1063/1.870745>, URL <http://aip.scitation.org/doi/10.1063/1.870745>.
- [66] S.J. Allfrey, R. Hatzky, *Comput. Phys. Comm.* 154 (2) (2003) 98–104, [http://dx.doi.org/10.1016/S0010-4655\(03\)00288-1](http://dx.doi.org/10.1016/S0010-4655(03)00288-1), URL <https://www.sciencedirect.com/science/article/pii/S0010465503002881?via%3Dihub>.

- [67] J.H. Halton, Numer. Math. 2 (1) (1960) 84–90, <http://dx.doi.org/10.1007/BF01386213>, URL <http://link.springer.com/10.1007/BF01386213>.
- [68] J.M. Hammersley, Ann. New York Acad. Sci. 86 (3) (1960) 844–874, <http://dx.doi.org/10.1111/j.1749-6632.1960.tb42846.x>, URL <http://doi.wiley.com/10.1111/j.1749-6632.1960.tb42846.x>.
- [69] J. van der Corput, Proc. Akad. Wet. Amsterdam 38 (1935) 813–821.
- [70] B. McMillan, S. Jolliet, A. Bottino, P. Angelino, T. Tran, L. Villard, Comput. Phys. Comm. 181 (4) (2010) 715–719, <http://dx.doi.org/10.1016/j.cpc.2009.12.001>, URL <http://linkinghub.elsevier.com/retrieve/pii/S0010465509003920> <http://www.sciencedirect.com/science/article/pii/S0010465509003920>.
- [71] M. Frigo, S.G. Johnson, Proc. IEEE 93 (2) (2005) 216–231, <http://dx.doi.org/10.1109/JPROC.2004.840301>, URL <http://ieeexplore.ieee.org/document/1386650/>.
- [72] E. Anderson, Z. Bai, C. Bischof, L.S. Blackford, J. Demmel, J. Dongarra, J. Du Croz, A. Greenbaum, S. Hammarling, A. McKenney, D. Sorensen, LAPACK Users' Guide, Society for Industrial and Applied Mathematics, 1999, <http://dx.doi.org/10.1137/1.9780898719604>, URL <http://epubs.siam.org/doi/book/10.1137/1.9780898719604>.
- [73] R. Hatzky, T.M. Tran, A. Könies, R. Kleiber, S.J. Allfrey, Phys. Plasmas 9 (3) (2002) 898, <http://dx.doi.org/10.1063/1.1449889>, URL <http://aip.scitation.org/doi/10.1063/1.1449889>.
- [74] M.N. Rosenbluth, F.L. Hinton, Phys. Rev. Lett. 80 (4) (1998) 724–727, <http://dx.doi.org/10.1103/PhysRevLett.80.724>, URL <http://link.aps.org/doi/10.1103/PhysRevLett.80.724>.
- [75] Y. Chen, S.E. Parker, Phys. Plasmas 14 (8) (2007) 082301, <http://dx.doi.org/10.1063/1.2751603>, URL <http://aip.scitation.org/doi/10.1063/1.2751603>.
- [76] W. Dorland, Phys. Plasmas 8 (5) (2001) 2095–2100, <http://dx.doi.org/10.1063/1.1351828>, URL <http://aip.scitation.org/doi/10.1063/1.1351828>.
- [77] Y. Chen, S.E. Parker, J. Comput. Phys. 189 (2) (2003) 463–475, [http://dx.doi.org/10.1016/S0021-9991\(03\)00228-6](http://dx.doi.org/10.1016/S0021-9991(03)00228-6), URL <https://www.sciencedirect.com/science/article/pii/S0021999103002286>.
- [78] A. Mishchenko, R. Hatzky, A. Könies, Phys. Plasmas 11 (12) (2004) 5480–5486, <http://dx.doi.org/10.1063/1.1812275>, URL <http://aip.scitation.org/doi/10.1063/1.1812275>.
- [79] A. Mishchenko, A. Könies, R. Hatzky, Proc. Jt. Varenna-Lausanne Int. Work., Società Italiana di Fisica, Bologna, 2004, p. 315.
- [80] A. Mishchenko, A. Könies, R. Kleiber, M. Cole, Phys. Plasmas 21 (9) (2014) 092110, <http://dx.doi.org/10.1063/1.4895501>, arXiv:1407.3992. URL <http://aip.scitation.org/doi/10.1063/1.4895501>.
- [81] R. Kleiber, R. Hatzky, A. Könies, A. Mishchenko, E. Sonnendrücker, Phys. Plasmas 23 (3) (2016) 032501, <http://dx.doi.org/10.1063/1.4942788>, URL <http://aip.scitation.org/doi/10.1063/1.4942788>.
- [82] A. Mishchenko, M. Cole, R. Kleiber, A. Könies, Phys. Plasmas 21 (5) (2014) 052113, <http://dx.doi.org/10.1063/1.4880560>, URL <http://aip.scitation.org/doi/10.1063/1.4880560>.
- [83] A. Biancalani, A. Bottino, S. Briguglio, A. Könies, P. Lauber, A. Mishchenko, E. Poli, B.D. Scott, F. Zonca, Phys. Plasmas 23 (1) (2016) 012108, <http://dx.doi.org/10.1063/1.4939803>, arXiv:1510.01945. URL <http://aip.scitation.org/doi/10.1063/1.4939803>.
- [84] A. Biancalani, A. Bottino, M. Cole, C. Di Troia, P. Lauber, A. Mishchenko, B. Scott, F. Zonca, Plasma Phys. Control. Fusion 59 (5) (2017) 054004, <http://dx.doi.org/10.1088/1361-6587/aa61e4>, arXiv:1610.01960. URL <http://stacks.iop.org/0741-3335/59/i=5/a=054004?key=crossref.5535f60889456ea6eafc60d759965eb5>.
- [85] P. Catto, W. Tang, D. Baldwin, Plasma Phys. 23 (7) (1981) 639–650, <http://dx.doi.org/10.1088/0032-1028/23/7/005>, URL <http://stacks.iop.org/0032-1028/23/i=7/a=005?key=crossref.ec4e34417bc78b7c7d606d8fa458b827>.
- [86] B.F. McMillan, L. Villard, Phys. Plasmas 21 (5) (2014) 052501, <http://dx.doi.org/10.1063/1.4873387>, URL <http://aip.scitation.org/doi/10.1063/1.4873387>.
- [87] C.C. Kim, S.E. Parker, J. Comput. Phys. 161 (2) (2000) 589–604, <http://dx.doi.org/10.1006/jcph.2000.6518>, URL <https://www.sciencedirect.com/science/article/pii/S0021999100965185>.
- [88] R. Hatzky, Parallel Comput. 32 (4) (2006) 325–330, <http://dx.doi.org/10.1016/j.parco.2006.03.001>, URL <https://www.sciencedirect.com/science/article/pii/S0167819106000111>.
- [89] N. Ohana, C. Gheller, E. Lanti, A. Jocksch, S. Brunner, L. Villard, 2019. Gyrokinetic Simulations on Many- and Multi-core Architectures with the Global Electromagnetic Particle-In-Cell Code ORB5, aug 2019, eprint 1908.02219, arXiv:1908.02219.
- [90] A.G. Peeters, C. Angioni, Y. Camenen, F.J. Casson, W.A. Hornsby, A.P. Snodin, D. Strintzi, Phys. Plasmas 16 (6) (2009) 062311, <http://dx.doi.org/10.1063/1.3124133>, URL <http://aip.scitation.org/doi/10.1063/1.3124133>.
- [91] M.D.J. Cole, A. Biancalani, A. Bottino, R. Kleiber, A. Könies, A. Mishchenko, Phys. Plasmas 24 (2) (2017) 022508, <http://dx.doi.org/10.1063/1.4976322>, URL <http://aip.scitation.org/doi/10.1063/1.4976322>.
- [92] P. Angelino, X. Garbet, L. Villard, A. Bottino, S. Jolliet, P. Ghendrih, V. Grandgirard, B.F. McMillan, Y. Sarazin, G. Dif-Pradalier, T.M. Tran, Phys. Plasmas 15 (6) (2008) 062306, <http://dx.doi.org/10.1063/1.2928849>, URL <http://aip.scitation.org/doi/10.1063/1.2928849>.
- [93] A. Biancalani, A. Bottino, S. Brunner, A.D. Siena, T. Görler, R. Hatzky, F. Jenko, A. Könies, P. Lauber, I. Novikau, A. Mishchenko, F. Vannini, L. Villard, A. Zocco, 46thEPS Conf. Plasma Phys., 2019, URL <http://ocs.ciemat.es/EPS2019ABS/pdf/15.J602.pdf>.
- [94] A. Biancalani, A. Bottino, F. Jenko, P. Lauber, I. Novikau, A. Mishchenko, L. Villard, Submitted to Phys. Rev. Lett. (2019).